

GENERATORS OF INVARIANTS OF TWO 4×4 MATRICES

VESSELIN DRENSKY AND LILIYA SADIKOVA

ABSTRACT. Over a field of characteristic 0, the algebra of invariants of several $n \times n$ matrices under simultaneous conjugation by GL_n is generated by traces of products of generic matrices. In this paper we have found, in terms of representation theory of GL_2 , a minimal set of generators of the algebra of invariants of two 4×4 matrices. The proof is purely combinatorial and involves computer calculations with standard functions of Maple.

INTRODUCTION

Let K be a field of characteristic 0 and let C_{nd} be the pure trace algebra generated by all traces of products $\text{tr}(X_{i_1} \cdots X_{i_k})$ of the generic $n \times n$ matrices $X_i = (x_{pq}^{(i)})$, $p, q = 1, \dots, n$, $i = 1, \dots, d$. The algebra C_{nd} coincides with the algebra of invariants of the general linear group $GL_n = GL_n(K)$ acting by simultaneous conjugation on d matrices of size $n \times n$.

For a background on the algebras of matrix invariants see, e.g. [F, DF]. Traditionally, a result giving the explicit generators of the algebra of invariants of a linear group G is called a first fundamental theorem of the invariant theory of G and a result describing the relations between the generators a second fundamental theorem. Classical invariant theory implies that C_{nd} is finitely generated. By the Noether normalization theorem C_{nd} contains a homogeneous set of algebraically independent elements $\{f_1, \dots, f_k\}$ called a homogeneous system of parameters, where $k = (d-1)n^2 + 1$ is the transcendence degree of the quotient field of C_{nd} , such that C_{nd} is integral over the polynomial algebra $S = K[f_1, \dots, f_k]$. A more precise result, see Van den Bergh [V], gives that C_{nd} is a finitely generated free module over a suitable chosen S . An upper bound for the generating set of the algebra C_{nd} is given in terms of PI-algebras. By the Nagata-Higman theorem the polynomial identity $x^n = 0$ implies the identity $x_1 \cdots x_N = 0$ for some $N = N(n)$. If N is minimal with this property, then C_{nd} is generated by traces of products $\text{tr}(X_{i_1} \cdots X_{i_k})$ of degree $k \leq N$. This estimate is sharp if d is sufficiently large. The values of $N(n)$ are between $n(n+1)/2$ and n^2 , bounds obtained respectively by Kuzmin [Ku] and Razmyslov [R] (see also [DF] for an exposition of the results in [Ku]). The only explicitly known values of $N(n)$ are $N(2) = 3$ and $N(3) = 6$ (an old result of Dubnov [Du]), and $N(4) = 10$ (a result of Vaughan-Lee [VL]). Recently, Shestakov and Zhukavets [SZ] obtained that the class of nilpotency of 2-generated nil algebras of nil index 5 is 15. All these results agree with the conjecture of Kuzmin [Ku] that $N(n) = n(n+1)/2$.

2000 *Mathematics Subject Classification.* 16R30.

Key words and phrases. Matrix invariants, trace algebras.

The first author was partially supported by Grant MM1106/2001 of the Bulgarian National Science Fund.

Explicit minimal sets of generators of C_{nd} are found in few cases only. By a theorem of Sibirskii [S], C_{2d} is generated by $\text{tr}(X_i)$, $1 \leq i \leq d$, $\text{tr}(X_i X_j)$, $1 \leq i \leq j \leq d$, $\text{tr}(X_i X_j X_k)$, $1 \leq i < j < k \leq d$. Teranishi [T1] found the following system of generators of C_{32} :

$$\begin{aligned} &\text{tr}(X), \text{tr}(Y), \text{tr}(X^2), \text{tr}(XY), \text{tr}(Y^2), \\ &\text{tr}(X^3), \text{tr}(X^2Y), \text{tr}(XY^2), \text{tr}(Y^3), \text{tr}(X^2Y^2), \text{tr}(X^2Y^2XY), \end{aligned}$$

where X, Y are generic 3×3 matrices. He showed that the first ten of these generators form a homogeneous system of parameters of C_{32} and C_{32} is a free module with generators 1 and $\text{tr}(X^2Y^2XY)$ over the polynomial algebra on these ten elements. Abeasis and Pittaluga [AP] found a system of generators of C_{3d} , for any $d \geq 2$, in terms of representation theory of the symmetric and general linear groups, in the spirit of its use in theory of PI-algebras. Teranishi [T1, T2] found also a homogeneous system of parameters of C_{42} and a set of generators, all of them being traces of products of generic matrices.

It has turned out that the systems of generators of Sibirskii [S] for C_{2d} and of Teranishi [T1] for C_{32} are not very convenient if we want to find the defining relations of the algebra. (Compare the relation for C_{23} found by Sibirskii [S] with the defining relations for C_{2d} of one of the authors [D2] with respect to another natural system of generators.) Also, it follows from the description of the generators of C_{32} , that $\text{tr}(X^2Y^2XY)$ satisfies a quadratic equation with coefficients depending on the other ten generators. The explicit (but very complicated) form of the equation was found by Nakamoto [N], over $\mathbb{Z}$, with respect to a slightly different system of generators. A much simple relation with respect to the system of generators of C_{32} in [AP] was found by Aslaksen, Drensky and Sadikova in [ADS].

The purpose of the present paper is to find a minimal system of generators of C_{42} in the spirit of the work of Abeasis and Pittaluga [AP] used in their description of the generators of C_{nd} . Applied to C_{42} , the idea in [AP] is the following. The general linear group GL_2 acts canonically on the free associative algebra $K\langle x, y \rangle$ and this induces an action on the pure trace algebra

$$P(x, y) = K[\text{tr}(z_1 \cdots z_k) \mid z_i = x, y, \ i = 1, \dots, k, \ k = 1, 2, \dots],$$

and on the algebra C_{42} (because C_{42} is a homomorphic image of $P(x, y)$ under the natural homomorphism extending the map $x \rightarrow X, y \rightarrow Y$, where X and Y are two generic 4×4 matrices). The algebra C_{42} has a system of generators of degree ≤ 10 . Without loss of generality we may assume that this system consists of traces of products $\text{tr}(Z_1 \cdots Z_k)$, $Z_i = X, Y$. Let T_k be the subalgebra of C_{42} generated by all traces $\text{tr}(Z_1 \cdots Z_l)$ of degree $l \leq k$, $Z_i = X, Y$. Clearly, T_k is also a GL_2 -submodule of C_{42} . Let $C_{42}^{(k+1)}$ be the homogeneous component of degree $k+1$ of C_{42} . Then the intersection $T_k \cap C_{42}^{(k+1)}$ is a GL_2 -module and has a complement G_{k+1} in $C_{42}^{(k+1)}$, which is the GL_2 -module of the “new” generators of degree $k+1$. We may assume that G_{k+1} is a submodule of the GL_2 -module spanned by traces of products $\text{tr}(Z_1 \cdots Z_{k+1})$ of degree $k+1$. The GL_2 -module of the generators of C_{42} is

$$G = G_1 \oplus G_2 \oplus \cdots \oplus G_{10}.$$

The authors of [AP] work with the multilinear elements in the free algebra which are consequences of the polynomial identity $x^n = 0$. Instead, we prefer to work

directly in the algebras $P(x, y)$ and C_{42} with more essential use of representation theory of the general linear group, in the spirit of [D1]. Our main result is that the minimal generating GL_2 -module G of C_{42} decomposes as

$$G = W(1, 0) \oplus W(2, 0) \oplus W(3, 0) \oplus W(4, 0) \oplus W(2, 2) \oplus W(3, 2) \\ \oplus W(4, 2) \oplus W(3, 3) \oplus W(4, 3) \oplus W(5, 3) \oplus W(4, 4) \oplus W(6, 3) \oplus W(5, 5),$$

where $W(\lambda_1, \lambda_2)$ is the irreducible GL_2 -module corresponding to the partition (λ_1, λ_2) . For $\lambda = (\lambda_1, \lambda_2) \neq (5, 5)$, one may choose as a generator of $W(\lambda_1, \lambda_2)$ the element

$$w_\lambda(X, Y) = \text{tr}((XY - YX)^{\lambda_2} X^{\lambda_1 - \lambda_2}).$$

For $\lambda = (5, 5)$ a generator of $W(5, 5)$ is

$$w_{(5,5)}(X, Y) = \text{tr}((XY - YX)^3 (X^2 Y^2 - XY YX - YX XY + Y^2 X^2)).$$

It seems to us that our generators are more convenient than the usual ones, if we want to find a system of defining relations of T_{42} . Most of the computations are performed with Maple, using standard functions only. Some of the obtained relations and technical results may be used for concrete calculations with traces of generic 4×4 matrices.

1. PRELIMINARIES

The group GL_2 acts in a canonical way on the vector space with basis $\{x, y\}$ and this action induces a diagonal action on the free algebra $K\langle x, y \rangle$:

$$g(z_1 \cdots z_k) = g(z_1) \cdots g(z_k), \quad z_i = x, y, \quad g \in GL_2.$$

The action of GL_2 on $K\langle x, y \rangle$ induces an action on the pure trace algebra $P(x, y)$:

$$g(\text{tr}(z_1 \cdots z_k)) = \text{tr}(g(z_1) \cdots g(z_k)), \quad z_i = x, y, \quad g \in GL_2.$$

If we fix the generic $n \times n$ matrices X, Y , the algebra of invariants C_{n2} becomes a homomorphic image of $P(x, y)$ under the natural homomorphism extending the map $x \rightarrow X, y \rightarrow Y$ and the GL_2 action is transferred on C_{n2} . The GL_2 -module $K\langle x, y \rangle$ is a direct sum of irreducible polynomial modules, described in terms of partitions $\lambda = (\lambda_1, \lambda_2)$. We denote by $W(\lambda)$ the corresponding GL_2 -module.

The GL_2 -submodules and factor modules W of $K\langle x, y \rangle$ inherit its natural bi-grading which counts the entries of x and y in each monomial. We denote by $W^{(p,q)}$ the corresponding homogeneous component. The formal power series

$$H(W, t, u) = \sum_{p,q \geq 0} \dim(W^{(p,q)}) t^p u^q$$

is called the Hilbert series of W . The Hilbert series of $W(\lambda)$ is the Schur function $S_\lambda(t, u)$ which, in the case of two variables, has the very simple form

$$S_\lambda(t, u) = (tu)^{\lambda_2} (t^{\lambda_1 - \lambda_2} + t^{\lambda_1 - \lambda_2 - 1} u + \cdots + tu^{\lambda_1 - \lambda_2 - 1} + u^{\lambda_1 - \lambda_2}).$$

The Hilbert series of W plays the role of its character, and

$$W = \bigoplus W^{\oplus m(\lambda)}(\lambda), \quad m(\lambda) \in \mathbb{N} \cup \{0\},$$

i.e. $W(\lambda)$ participates in W with multiplicity $m(\lambda)$, if and only if

$$H(W, t, u) = \sum m(\lambda) S_\lambda(t, u).$$

The Hilbert series of C_{42} was calculated by Teranishi [T2], with some misprints in the expression, and, corrected, by Berele and Stembridge [BS]:

$$(1) \quad H(C_{42}, t, u) = \frac{P_C(t, u)}{(1-t)(1-u)Q_C(t, u)},$$

$$P_C(t, u) = (1 - e_2 + e_2^2)(1 - e_1e_2 + e_1e_2^2 + e_1^2e_2^2 + e_1e_2^3 - e_1e_2^4 + e_2^6),$$

$$e_1 = t + u, \quad e_2 = tu,$$

$$Q_C(t, u) = (1 - t^2)(1 - t^3)(1 - t^4)(1 - u^2)(1 - u^3)(1 - u^4)$$

$$(1 - tu)^2(1 - t^2u)^2(1 - tu^2)^2(1 - t^3u)(1 - tu^3)(1 - t^2u^2).$$

We shall need the decomposition as a sum of Schur functions of the first 11 homogeneous components $h_n(t, u)$ of the symmetric formal power series

$$(2) \quad h(t, u) = \frac{P_C(t, u)}{Q_C(t, u)} = (1-t)(1-u)H(C_{42}, t, u) = \sum_{n \geq 0} h_n(t, u).$$

Using the formula

$$(3) \quad \frac{1}{1-z} = 1 + z + z^2 + \dots,$$

direct calculations show that

$$(4) \quad \begin{aligned} h_0 &= 1, \quad h_1 = 0, \quad h_2 = t^2 + tu + u^2 = S_{(2,0)}(t, u), \\ h_3 &= S_{(3,0)}(t, u), \quad h_4 = 2S_{(4,0)}(t, u) + 2S_{(2,2)}(t, u), \\ h_5 &= S_{(5,0)}(t, u) + S_{(4,1)}(t, u) + 2S_{(3,2)}(t, u), \\ h_6 &= 3S_{(6,0)}(t, u) + S_{(5,1)}(t, u) + 5S_{(4,2)}(t, u) + S_{(3,3)}(t, u), \\ h_7 &= 2S_{(7,0)}(t, u) + 2S_{(6,1)}(t, u) + 5S_{(5,2)}(t, u) + 4S_{(4,3)}(t, u), \\ h_8 &= 4S_{(8,0)}(t, u) + 2S_{(7,1)}(t, u) + 10S_{(6,2)}(t, u) + 6S_{(5,3)}(t, u) + 8S_{(4,4)}(t, u), \\ h_9 &= 3S_{(9,0)}(t, u) + 3S_{(8,1)}(t, u) + 10S_{(7,2)}(t, u) + 13S_{(6,3)}(t, u) + 8S_{(5,4)}(t, u), \\ h_{10} &= 5S_{(10,0)}(t, u) + 4S_{(9,1)}(t, u) + 16S_{(8,2)}(t, u) \\ &\quad + 16S_{(7,3)}(t, u) + 24S_{(6,4)}(t, u) + 5S_{(5,5)}(t, u). \end{aligned}$$

Any submodule $W(\lambda) = W(\lambda_1, \lambda_2)$ of $K\langle x, y \rangle$ is generated by a unique, up to a multiplicative constant, homogeneous element $w_\lambda(x, y)$ of degree λ_1 and λ_2 with respect to x and y , respectively, called the highest weight vector of $W(\lambda)$. It is characterized by the following property, see [DEP, ADF] and [K] for the version which we need. We state it for two variables only. The homogeneous polynomial $w_\lambda(x, y) \in K\langle x, y \rangle$ of degree (λ_1, λ_2) is a highest weight vector for some $W(\lambda_1, \lambda_2)$ if and only if $\delta(w_\lambda(x, y)) = 0$, where δ is the derivation of $K\langle x, y \rangle$ defined by $\delta(x) = 0$, $\delta(y) = x$. If W_i , $i = 1, \dots, k$, are k isomorphic copies of $W(\lambda)$ and $w_i \in W_i$ are highest weight vectors, then the highest weight vector of any submodule $W(\lambda)$ of the direct sum $W_1 \oplus \dots \oplus W_k$ has the form $\xi_1 w_1 + \dots + \xi_k w_k$ for some $\xi_i \in K$. Any k linearly independent highest weight vectors can serve as generators of the GL_2 -module $W_1 \oplus \dots \oplus W_k$. Of course, similar arguments work for submodules and homomorphic images of the GL_2 -module $K\langle x, y \rangle$.

The multiplicity of $W(\lambda)$ in $K\langle x, y \rangle$ is equal to the degree $d(\lambda)$ of the corresponding irreducible representation of the symmetric group S_k , $k = \lambda_1 + \lambda_2$, or, equivalently, to the number of standard λ -tableaux. For the standard λ -tableau

$$T_\sigma = \begin{array}{|c|c|c|c|c|c|} \hline \sigma(1) & \cdots & \sigma(2\lambda_2 - 1) & \sigma(2\lambda_2 + 1) & \cdots & \sigma(k) \\ \hline \sigma(2) & \cdots & \sigma(2\lambda_2) & & & \\ \hline \end{array}$$

corresponding to $\sigma \in S_k$, $k = \lambda_1 + \lambda_2$, we associate a highest weight vector $w(T_\sigma)$ in $K\langle x, y \rangle$. When $\sigma = \varepsilon$ is the identity of S_k we fix

$$\begin{aligned} w(T_\varepsilon) &= (xy - yx)^{\lambda_2} x^{\lambda_1 - \lambda_2} \\ &= \sum_{\rho_1, \dots, \rho_s \in S_2} \text{sign}(\rho_1 \cdots \rho_s) z_{\rho_1(1)} z_{\rho_1(2)} \cdots z_{\rho_s(1)} z_{\rho_s(2)} x^{\lambda_1 - \lambda_2}, \end{aligned}$$

where $z_1 = x$, $z_2 = y$, $s = \lambda_2$. For σ arbitrary, we define $w(T_\sigma)$ in a similar way, but the skew-symmetries are in positions $(\sigma(1), \sigma(2)), \dots, (\sigma(2s-1), \sigma(2s))$ instead of the positions $(1, 2), \dots, (2s-1, 2s)$ (and the positions with fixed x are $\sigma(2s+1), \dots, \sigma(k)$ instead of $2s+1, \dots, k$). For the standard (p, q) -tableau

$$T = \begin{array}{|c|c|c|c|c|c|} \hline a_1 & \cdots & a_q & a_{q+1} & \cdots & a_p \\ \hline b_1 & \cdots & b_q & & & \\ \hline \end{array}$$

we denote the corresponding highest weight vector $\text{tr}(w(T))$ by

$$w(T) = w \begin{pmatrix} a_1 & \cdots & a_q & a_{q+1} & \cdots & a_p \\ b_1 & \cdots & b_q & & & \end{pmatrix}.$$

2. THE GL_2 -MODULES OF FORMAL TRACES

We denote by U_n the vector subspace of all traces of formal products of length n in the variables x and y in the pure trace algebra $P(x, y)$.

Proposition 2.1. *The GL_2 -modules U_n spanned by the traces of the products of length $n \leq 10$ are decomposed as follows:*

$$\begin{aligned} U_1 &= W(1), & U_2 &= W(2), & U_3 &= W(3), & U_4 &= W(4) \oplus W(2, 2), \\ U_5 &= W(5) \oplus W(3, 2), & U_6 &= W(6) \oplus 2W(4, 2) \oplus W(3, 3), \\ U_7 &= W(7) \oplus 2W(5, 2) \oplus 2W(4, 3), \\ U_8 &= W(8) \oplus 3W(6, 2) \oplus 3W(5, 3) \oplus 3W(4, 4), \\ U_9 &= W(9) \oplus 3W(7, 2) \oplus 6W(6, 3) \oplus 4W(5, 4), \\ U_{10} &= W(10) \oplus 4W(8, 2) \oplus 7W(7, 3) \oplus 10W(6, 4) \oplus 4W(5, 5). \end{aligned}$$

Proof. We calculate the Hilbert series $H(U_n, t, u)$ of U_n . Since $H(U_n, t, u)$ is a symmetric polynomial in t, u , it is sufficient to count the number h_{nk} of traces of degree k with respect to x and $n - k$ with respect to y , $k \geq n - k$. There is one trace depending on x only, $\text{tr}(x^n)$, and one trace, $\text{tr}(x^{n-1}y)$, of degree $(n-1, 1)$. Hence $h_{n0} = h_{n1} = 1$ and we may consider the case $k \leq n-2$ only. Since the traces are invariant under cyclic permutation, we may assume that they are of the form

$$\text{tr}(x^{a_1} y^{b_1} \cdots x^{a_p} y^{b_p}),$$

where $a_i, b_i > 0$ and $a_1 + \cdots + a_p = k$, $b_1 + \cdots + b_p = n - k$. We order the words $x^{a_1} y^{b_1} \cdots x^{a_p} y^{b_p}$ comparing lexicographically the $2p$ -tuples $(a_1, \dots, a_p, b_1, \dots, b_p)$ and fix the representative of $\text{tr}(x^{a_1} y^{b_1} \cdots x^{a_p} y^{b_p})$ to be as high as possible.

Cases $n = 1, 2, 3$. These cases are obvious because the coefficients $h_{nk}, h_{n, n-k}$ of $H(U_n, t, v)$ are equal to 1 for $k = 0, 1$ and

$$\begin{aligned} H(U_1, t, v) &= t + u = S_{(1)}(t, u), \\ H(U_2, t, u) &= t^2 + tu + u^2 = S_{(2)}(t, u), \\ H(U_3, t, u) &= t^3 + t^2u + tu^2 + u^3 = S_{(3)}(t, u). \end{aligned}$$

This means that $U_1 = W(1)$, $U_2 = W(2)$ and $U_3 = W(3)$.

Case $n = 4$. The only elements of degree $(2, 2)$ are $\text{tr}(x^2y^2)$, $\text{tr}(xyxy)$ and this implies

$$H(U_4, t, u) = t^4 + t^3u + 2t^2u^2 + tu^3 + u^4 = S_{(4)}(t, u) + S_{(2^2)}(t, u),$$

and hence $U_4 = W(4) \oplus W(2, 2)$.

Case $n = 5$. The traces of degree $(3, 2)$ are $\text{tr}(x^3y^2)$, $\text{tr}(x^2yxy)$ and

$$H(U_5, t, u) = t^5 + t^4u + 2t^3u^2 + 2t^2u^3 + tu^4 + u^5 = S_{(5)}(t, u) + S_{(3,2)}(t, u),$$

which gives $U_5 = W(5) \oplus W(3, 2)$.

Case $n = 6$. The traces of degree $(4, 2)$ are $\text{tr}(x^4y^2)$, $\text{tr}(x^3yxy)$, $\text{tr}(x^2yx^2y)$, these of degree $(3, 3)$ are $\text{tr}(x^3y^3)$, $\text{tr}(x^2y^2xy)$, $\text{tr}(x^2yxy^2)$, $\text{tr}(xyxyxy)$, and

$$\begin{aligned} H(U_6, t, u) &= t^6 + t^5u + 3t^4u^2 + 4t^3u^3 + 3t^2u^4 + tu^5 + u^6 \\ &= S_{(6)}(t, u) + 2S_{(4,2)}(t, u) + S_{(3^2)}(t, u), \\ U_6 &= W(6) \oplus 2W(4, 2) \oplus W(3, 3). \end{aligned}$$

Case $n = 7$. The traces of degree $(5, 2)$ are $\text{tr}(x^5y^2)$, $\text{tr}(x^4yxy)$, $\text{tr}(x^3yx^2y)$, these of degree $(4, 3)$ are $\text{tr}(x^4y^3)$, $\text{tr}(x^3y^2xy)$, $\text{tr}(x^3yxy^2)$, $\text{tr}(x^2y^2x^2y)$, $\text{tr}(x^2yxyxy)$ and

$$\begin{aligned} H(U_7, t, u) &= t^7 + t^6u + 3t^5u^2 + 5t^4u^3 + 5t^3u^4 + 3t^2u^5 + tu^6 + u^7 \\ &= S_{(7)}(t, u) + 2S_{(5,2)}(t, u) + 2S_{(4,3)}(t, u), \\ U_7 &= W(7) \oplus 2W(5, 2) \oplus 2W(4, 3). \end{aligned}$$

Case $n = 8$. The traces of degree $(6, 2)$ are

$$\text{tr}(x^6y^2), \quad \text{tr}(x^{a_1}yx^{a_2}y), \quad (a_1, a_2) = (5, 1), (4, 2), (3, 3),$$

and $h_{82} = 4$, the traces of degree $(5, 3)$ are

$$\text{tr}(x^5y^3),$$

$$\begin{aligned} \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \quad (a_1, a_2) = (4, 1), (3, 2), \quad (b_1, b_2) = (2, 1), (1, 2), \\ \text{tr}(x^{a_1}yx^{a_2}yx^{a_3}y), \quad (a_1, a_2, a_3) = (3, 1, 1), (2, 2, 1), \end{aligned}$$

$h_{83} = 7$. Finally, the traces of degree $(4, 4)$ are

$$\begin{aligned} \text{tr}(x^4y^4), \quad \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \\ (a_1, a_2) = (3, 1), (b_1, b_2) = (3, 1), (2, 2), (1, 3), \\ (a_1, a_2) = (2, 2), (b_1, b_2) = (3, 1), (2, 2), \\ \text{tr}(x^2y^{b_1}xy^{b_2}xy^{b_3}), \quad (b_1, b_2, b_3) = (2, 1, 1), (1, 2, 1), (1, 1, 2), \\ \text{tr}(xyxyxyxy). \end{aligned}$$

Hence $h_{84} = 10$,

$$\begin{aligned} H(U_8, t, u) &= t^8 + t^7u + 4t^6u^2 + 7t^5u^3 + 10t^4u^4 + 7t^3u^5 + 4t^2u^6 + tu^7 + u^8 \\ &= S_{(8)}(t, u) + 3S_{(6,2)}(t, u) + 3S_{(5,3)}(t, u) + 3S_{(4,4)}(t, u), \\ U_8 &= W(8) \oplus 3W(6, 2) \oplus 3W(5, 3) \oplus 3W(4, 4). \end{aligned}$$

Case $n = 9$. The traces of degree $(7, 2)$ are

$$\text{tr}(x^7y^2), \quad \text{tr}(x^{a_1}yx^{a_2}y), \quad (a_1, a_2) = (6, 1), (5, 2), (4, 3),$$

and $h_{92} = 4$, these of degree $(6, 3)$ are

$$\begin{aligned} \text{tr}(x^6y^3), \quad \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \\ (a_1, a_2) = (5, 1), (4, 2), (b_1, b_2) = (2, 1), (1, 2), \quad (a_1, a_2) = (3, 3), (b_1, b_2) = (2, 1), \end{aligned}$$

$\text{tr}(x^{a_1}yx^{a_2}yx^{a_3}y)$, $(a_1, a_2, a_3) = (4, 1, 1), (3, 2, 1), (3, 1, 2), (2, 2, 2)$,
 $h_{93} = 10$. The traces of degree $(5, 4)$ are
 $\text{tr}(x^5y^4)$, $\text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2})$, $(a_1, a_2) = (4, 1), (3, 2), (b_1, b_2) = (3, 1), (2, 2), (1, 3)$,
 $\text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}x^{a_3}y^{b_3})$,
 $(a_1, a_2, a_3) = (3, 1, 1), (2, 2, 1)$, $(b_1, b_2, b_3) = (2, 1, 1), (1, 2, 1), (1, 1, 2)$,
 $\text{tr}(x^2yxyxyxy)$,

$h_{94} = 14$. The Hilbert series and the decomposition of U_9 are, respectively,

$$\begin{aligned} H(U_9, t, u) &= t^9 + t^8u + 4t^7u^2 + 10t^6u^3 + 14t^5u^4 \\ &\quad + 14t^4u^5 + 10t^3u^6 + 4t^2u^7 + tu^8 + u^9 \\ &= S_{(9)}(t, u) + 3S_{(7,2)}(t, u) + 6S_{(6,3)}(t, u) + 4S_{(5,4)}(t, u), \\ U_9 &= W(9) \oplus 3W(7, 2) \oplus 6W(6, 3) \oplus 4W(5, 4). \end{aligned}$$

Case $n = 10$. The traces of degree $(8, 2)$ are

$$\text{tr}(x^8y^2), \quad \text{tr}(x^{a_1}yx^{a_2}y), \quad (a_1, a_2) = (7, 1), (6, 2), (5, 3), (4, 4),$$

and $h_{10,2} = 5$, these of degree $(7, 3)$ are

$$\begin{aligned} &\text{tr}(x^7y^3), \quad \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \\ &(a_1, a_2) = (6, 1), (5, 2), (4, 3), \quad (b_1, b_2) = (2, 1), (1, 2), \\ &\text{tr}(x^{a_1}yx^{a_2}yx^{a_3}y), \quad (a_1, a_2, a_3) = (5, 1, 1), (4, 2, 1), (4, 1, 2), (3, 3, 1), (3, 2, 2), \end{aligned}$$

$h_{10,3} = 12$. The traces of degree $(6, 4)$ are

$$\begin{aligned} &\text{tr}(x^6y^4), \quad \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \quad (a_1, a_2) = (5, 1), (4, 2), (b_1, b_2) = (3, 1), (2, 2), (1, 3), \\ &(a_1, a_2) = (3, 3), (b_1, b_2) = (3, 1), (2, 2), \\ &\text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}x^{a_3}y^{b_3}), \\ &(a_1, a_2, a_3) = (4, 1, 1), (3, 2, 1), (3, 1, 2), (b_1, b_2, b_3) = (2, 1, 1), (1, 2, 1), (1, 1, 2), \\ &(a_1, a_2, a_3) = (2, 2, 2), (b_1, b_2, b_3) = (2, 1, 1), \\ &\text{tr}(x^{a_1}yx^{a_2}yx^{a_3}yx^{a_4}y), \quad (a_1, a_2, a_3, a_4) = (3, 1, 1, 1), (2, 2, 1, 1), (2, 1, 2, 1), \end{aligned}$$

$h_{10,4} = 22$. Finally, the traces of degree $(5, 5)$ are

$$\begin{aligned} &\text{tr}(x^5y^5), \quad \text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}), \\ &(a_1, a_2) = (4, 1), (3, 2), \quad (b_1, b_2) = (4, 1), (3, 2), (2, 3), (1, 4), \\ &\text{tr}(x^{a_1}y^{b_1}x^{a_2}y^{b_2}x^{a_3}y^{b_3}), \\ &(a_1, a_2, a_3) = (3, 1, 1), (2, 2, 1), \\ &(b_1, b_2, b_3) = (3, 1, 1), (1, 3, 1), (1, 1, 3), (2, 2, 1), (2, 1, 2), (1, 2, 2), \\ &\text{tr}(x^2y^{b_1}xy^{b_2}xy^{b_3}xy^{b_4}), \\ &(b_1, b_2, b_3, b_4) = (2, 1, 1, 1), (1, 2, 1, 1), (1, 1, 2, 1), (1, 1, 1, 2), \\ &\text{tr}(xyxyxyxyxy), \end{aligned}$$

hence $h_{10,5} = 26$. The Hilbert series and the decomposition of U_{10} are, respectively,

$$\begin{aligned} H(U_{10}, t, u) &= t^{10} + t^9u + 5t^8u^2 + 12t^7u^3 + 22t^6u^4 \\ &\quad + 26t^5u^5 + 22t^4u^6 + 12t^3u^7 + 5t^2u^8 + tu^9 + u^{10}, \\ &= S_{(10)}(t, u) + 4S_{(8,2)}(t, u) + 7S_{(7,3)}(t, u) + 10S_{(6,4)}(t, u) + 4S_{(5,5)}(t, u), \\ U_{10} &= W(10) \oplus 4W(8, 2) \oplus 7W(7, 3) \oplus 10W(6, 4) \oplus 4W(5, 5). \end{aligned}$$

□

Lemma 2.2. *For $\lambda = (\lambda_1, \lambda_2) \vdash n$, $n \leq 10$, the following trace polynomials form a basis of the subspace of U_n consisting of all λ -highest weight vectors:*

$\lambda = (n)$, $n \geq 1$:

$$w = w(1 \ \cdots \ n) = \text{tr}(x^n);$$

$\lambda = (2, 2)$:

$$w = \frac{1}{2}w\begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix} = \frac{1}{2}\text{tr}([x, y]^2) = -\text{tr}(x^2y^2) + \text{tr}(xyxy);$$

$\lambda = (3, 2)$:

$$w = w\begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 \end{pmatrix} = \text{tr}([x, y]^2x) = -\text{tr}(x^3y^2) + \text{tr}(x^2yxy);$$

$\lambda = (4, 2)$:

$$w_1 = w\begin{pmatrix} 1 & 3 & 5 & 6 \\ 2 & 4 \end{pmatrix} = \text{tr}([x, y]^2x^2) = -\text{tr}(x^4y^2) + 2\text{tr}(x^3yxy) - \text{tr}(x^2yx^2y),$$

$$w_2 = w\begin{pmatrix} 1 & 2 & 5 & 6 \\ 3 & 4 \end{pmatrix} = \text{tr}(x^4y^2) - \text{tr}(x^2yx^2y);$$

$\lambda = (3, 3)$:

$$w = \frac{1}{3}w\begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{pmatrix} = \text{tr}(x^2y^2xy) - \text{tr}(x^2yxy^2);$$

$\lambda = (5, 2)$:

$$w_1 = w\begin{pmatrix} 1 & 3 & 5 & 6 & 7 \\ 2 & 4 \end{pmatrix} = \text{tr}([x, y]^2x^3) = -\text{tr}(x^5y^2) + 2\text{tr}(x^4yxy) - \text{tr}(x^3yx^2y),$$

$$w_2 = w\begin{pmatrix} 1 & 2 & 5 & 6 & 7 \\ 3 & 4 \end{pmatrix} = \text{tr}(x^5y^2) - \text{tr}(x^3yx^2y);$$

$\lambda = (4, 3)$:

$$w_1 = w\begin{pmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 \end{pmatrix} = \text{tr}(x^3y^2xy) - \text{tr}(x^3yxy^2),$$

$$w_2 = w\begin{pmatrix} 1 & 2 & 5 & 7 \\ 3 & 4 & 6 \end{pmatrix} = -\text{tr}(x^4y^3) + 2\text{tr}(x^3y^2xy) \\ + \text{tr}(x^3yxy^2) - \text{tr}(x^2y^2x^2y) - \text{tr}(x^2yxyxy);$$

$\lambda = (6, 2)$:

$$w_1 = w\begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 8 \\ 2 & 4 \end{pmatrix}$$

$$= \text{tr}([x, y]^2x^4) = -\text{tr}(x^6y^2) + 2\text{tr}(x^5yxy) - \text{tr}(x^4yx^2y),$$

$$w_2 = w\begin{pmatrix} 1 & 2 & 5 & 6 & 7 & 8 \\ 3 & 4 \end{pmatrix} = \text{tr}(x^6y^2) - \text{tr}(x^4yx^2y),$$

$$w_3 = w\begin{pmatrix} 1 & 3 & 4 & 6 & 7 & 8 \\ 2 & 5 \end{pmatrix} = -\text{tr}(x^6y^2) + \text{tr}(x^5yxy) + \text{tr}(x^4yx^2y) - \text{tr}(x^3yx^3y);$$

$\lambda = (5, 3)$:

$$w_1 = w\begin{pmatrix} 1 & 3 & 5 & 7 & 8 \\ 2 & 4 & 6 \end{pmatrix}$$

$$= \text{tr}(x^4y^2xy) - \text{tr}(x^4yxy^2) - \text{tr}(x^3y^2x^2y) + \text{tr}(x^3yx^2y^2),$$

$$w_2 = w\begin{pmatrix} 1 & 2 & 5 & 7 & 8 \\ 3 & 4 & 6 \end{pmatrix} = -\text{tr}(x^5y^3) + 2\text{tr}(x^4y^2xy)$$

$$-2\text{tr}(x^3y^2x^2y) + 2\text{tr}(x^3yx^2y^2) - \text{tr}(x^2yx^2yxy),$$

$$w_3 = w \begin{pmatrix} 1 & 3 & 5 & 6 & 8 \\ 2 & 4 & 7 & & \end{pmatrix} = \text{tr}(x^4y^2xy) - \text{tr}(x^4yxy^2);$$

$\lambda = (4, 4)$:

$$w_1 = \frac{1}{2}w \begin{pmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{pmatrix} = \frac{1}{2}\text{tr}([x, y]^4) = -2\text{tr}(x^2y^2xyxy) + 2\text{tr}(x^2yxy^2xy)$$

$$-2\text{tr}(x^2yxyxy^2) + \text{tr}(x^2y^2x^2y^2) + \text{tr}(xyxyxyxy),$$

$$w_2 = w \begin{pmatrix} 1 & 2 & 5 & 7 \\ 3 & 4 & 6 & 8 \end{pmatrix} = \text{tr}(x^3y^3xy) - 2\text{tr}(x^3y^2xy^2) + \text{tr}(x^3yxy^3)$$

$$- \text{tr}(x^2y^2xyxy) + 2\text{tr}(x^2yxy^2xy) - \text{tr}(x^2yxyxy^2) - 2\text{tr}(x^2y^3x^2y) + 2\text{tr}(x^2y^2x^2y^2),$$

$$w_3 = \frac{1}{2}w \begin{pmatrix} 1 & 2 & 5 & 6 \\ 3 & 4 & 7 & 8 \end{pmatrix} = \text{tr}(x^4y^4) - 2\text{tr}(x^3y^2xy^2) + \text{tr}(x^2yxy^2xy)$$

$$- 2\text{tr}(x^2y^3x^2y) + 2\text{tr}(x^2y^2x^2y^2);$$

$\lambda = (7, 2)$:

$$w_1 = w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 8 & 9 \\ 2 & 4 & & & & & \end{pmatrix}$$

$$= \text{tr}([x, y]^2x^5) = -\text{tr}(x^7y^2) + 2\text{tr}(x^6yxy) - \text{tr}(x^5yx^2y),$$

$$w_2 = w \begin{pmatrix} 1 & 2 & 5 & 6 & 7 & 8 & 9 \\ 3 & 4 & & & & & \end{pmatrix} = \text{tr}(x^7y^2) - \text{tr}(x^5yx^2y),$$

$$w_3 = w \begin{pmatrix} 1 & 2 & 4 & 6 & 7 & 8 & 9 \\ 3 & 5 & & & & & \end{pmatrix} = \text{tr}(x^6yxy) - \text{tr}(x^4yx^3y);$$

$\lambda = (6, 3)$:

$$w_1 = w \begin{pmatrix} 1 & 3 & 5 & 7 & 8 & 9 \\ 2 & 4 & 6 & & & \end{pmatrix} = \text{tr}([x, y]^3x^3) = \text{tr}(x^5y^2xy) - \text{tr}(x^5yxy^2)$$

$$- \text{tr}(x^4y^2x^2y) + \text{tr}(x^4yx^2y^2) - \text{tr}(x^3yx^2yxy) + \text{tr}(x^3yxyx^2y),$$

$$w_2 = w \begin{pmatrix} 1 & 2 & 5 & 7 & 8 & 9 \\ 3 & 4 & 6 & & & \end{pmatrix} = -\text{tr}(x^6y^3) + 2\text{tr}(x^5y^2xy)$$

$$- 2\text{tr}(x^4y^2x^2y) + \text{tr}(x^4yx^2y^2) + \text{tr}(x^3y^2x^3y) - \text{tr}(x^3yx^2yxy),$$

$$w_3 = w \begin{pmatrix} 1 & 2 & 4 & 7 & 8 & 9 \\ 3 & 5 & 6 & & & \end{pmatrix}$$

$$= \text{tr}(x^5yxy^2) - \text{tr}(x^5y^2xy) + \text{tr}(x^4yx^2y^2) - \text{tr}(x^4y^2x^2y),$$

$$w_4 = w \begin{pmatrix} 1 & 3 & 5 & 6 & 8 & 9 \\ 2 & 4 & 7 & & & \end{pmatrix} = \text{tr}(x^5y^2xy) - \text{tr}(x^5yxy^2) + \text{tr}(x^4yx^2y^2)$$

$$- \text{tr}(x^4yxyxy) - \text{tr}(x^3y^2x^3y) + \text{tr}(x^3yx^2yxy) + \text{tr}(x^3yxyx^2y) - \text{tr}(x^2yx^2yx^2y),$$

$$w_5 = w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 9 \\ 2 & 4 & 8 & & & \end{pmatrix}$$

$$= \text{tr}(x^5y^2xy) - \text{tr}(x^5yxy^2) + \text{tr}(x^3yx^2yxy) - \text{tr}(x^3yxyx^2y),$$

$$w_6 = w \begin{pmatrix} 1 & 2 & 4 & 6 & 8 & 9 \\ 3 & 5 & 7 & & & \end{pmatrix} = -\text{tr}(x^5y^2xy) + 2\text{tr}(x^4yx^2y^2)$$

$$+ \text{tr}(x^4yxyxy) - \text{tr}(x^3y^2x^3y) - \text{tr}(x^3yxyx^2y);$$

$\lambda = (5, 4)$:

$$\begin{aligned}
w_1 &= w \begin{pmatrix} 1 & 3 & 5 & 7 & 9 \\ 2 & 4 & 6 & 8 & \end{pmatrix} = \text{tr}(x^3 y^2 x^2 y^2) - \text{tr}(x^3 y^2 x y x y) \\
&+ \text{tr}(x^3 y x y^2 x y) - \text{tr}(x^3 y x y x y^2) - \text{tr}(x^2 y^2 x^2 y x y) + \text{tr}(x^2 y x y x y x y), \\
w_2 &= w \begin{pmatrix} 1 & 2 & 5 & 7 & 9 \\ 3 & 4 & 6 & 8 & \end{pmatrix} = \text{tr}(x^4 y^3 x y) - \text{tr}(x^4 y^2 x y^2) \\
&- \text{tr}(x^3 y^3 x^2 y) + \text{tr}(x^3 y x y x y^2) + \text{tr}(x^2 y^2 x^2 y x y) - \text{tr}(x^2 y x y x y x y), \\
w_3 &= w \begin{pmatrix} 1 & 2 & 4 & 7 & 9 \\ 3 & 5 & 6 & 8 & \end{pmatrix} = \text{tr}(x^4 y^3 x y) - \text{tr}(x^4 y x y^3) \\
&- \text{tr}(x^3 y^3 x^2 y) + \text{tr}(x^3 y x^2 y^3) - \text{tr}(x^3 y^2 x y x y) + \text{tr}(x^3 y x y x y^2), \\
w_4 &= w \begin{pmatrix} 1 & 2 & 3 & 4 & 9 \\ 5 & 6 & 7 & 8 & \end{pmatrix} = \text{tr}(x^5 y^4) - \text{tr}(x^4 y^3 x y) \\
&+ \text{tr}(x^4 y^2 x y^2) - \text{tr}(x^4 y x y^3) - \text{tr}(x^3 y x y^2 x y) + \text{tr}(x^2 y x y x y x y);
\end{aligned}$$

$\lambda = (8, 2)$:

$$\begin{aligned}
w_1 &= w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 8 & 9 & 10 \\ 2 & 4 & & & & & & \end{pmatrix} = -\text{tr}(x^8 y^2) + 2\text{tr}(x^7 y x y) - \text{tr}(x^6 y x^2 y), \\
w_2 &= w \begin{pmatrix} 1 & 3 & 4 & 6 & 7 & 8 & 9 & 10 \\ 2 & 5 & & & & & & \end{pmatrix} \\
&= -\text{tr}(x^8 y^2) + \text{tr}(x^7 y x y) + \text{tr}(x^6 y x^2 y) - \text{tr}(x^5 y x^3 y), \\
w_3 &= w \begin{pmatrix} 1 & 3 & 4 & 5 & 7 & 8 & 9 & 10 \\ 2 & 6 & & & & & & \end{pmatrix} \\
&= -\text{tr}(x^8 y^2) + \text{tr}(x^7 y x y) + \text{tr}(x^5 y x^3 y) - \text{tr}(x^4 y x^4 y), \\
w_4 &= w \begin{pmatrix} 1 & 3 & 4 & 5 & 6 & 8 & 9 & 10 \\ 2 & 7 & & & & & & \end{pmatrix} \\
&= -\text{tr}(x^8 y^2) + \text{tr}(x^7 y x y) - \text{tr}(x^5 y x^3 y) + \text{tr}(x^4 y x^4 y);
\end{aligned}$$

$\lambda = (7, 3)$:

$$\begin{aligned}
w_1 &= w \begin{pmatrix} 1 & 3 & 5 & 7 & 8 & 9 & 10 \\ 2 & 4 & 6 & & & & \end{pmatrix}, \\
w_2 &= w \begin{pmatrix} 1 & 3 & 5 & 6 & 8 & 9 & 10 \\ 2 & 4 & 7 & & & & \end{pmatrix}, \\
w_3 &= w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 9 & 10 \\ 2 & 4 & 8 & & & & \end{pmatrix}, \\
w_4 &= w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 & 8 & 10 \\ 2 & 4 & 9 & & & & \end{pmatrix}, \\
w_5 &= w \begin{pmatrix} 1 & 2 & 5 & 6 & 7 & 9 & 10 \\ 3 & 4 & 8 & & & & \end{pmatrix}, \\
w_6 &= w \begin{pmatrix} 1 & 3 & 4 & 7 & 8 & 9 & 10 \\ 2 & 5 & 6 & & & & \end{pmatrix}, \\
w_7 &= w \begin{pmatrix} 1 & 2 & 3 & 7 & 8 & 9 & 10 \\ 4 & 5 & 6 & & & & \end{pmatrix};
\end{aligned}$$

$\lambda = (6, 4)$:

$$w_1 = w \begin{pmatrix} 1 & 3 & 5 & 7 & 9 & 10 \\ 2 & 4 & 6 & 8 & & \end{pmatrix},$$

$$w_2 = w \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 7 & 8 & 9 & 10 & & \end{pmatrix},$$

$$w_3 = w \begin{pmatrix} 1 & 2 & 3 & 7 & 9 & 10 \\ 4 & 5 & 6 & 8 & & \end{pmatrix},$$

$$w_4 = w \begin{pmatrix} 1 & 2 & 5 & 7 & 9 & 10 \\ 3 & 4 & 6 & 8 & & \end{pmatrix},$$

$$w_5 = w \begin{pmatrix} 1 & 2 & 5 & 6 & 9 & 10 \\ 3 & 4 & 7 & 8 & & \end{pmatrix},$$

$$w_6 = w \begin{pmatrix} 1 & 2 & 5 & 6 & 7 & 10 \\ 3 & 4 & 8 & 9 & & \end{pmatrix},$$

$$w_7 = w \begin{pmatrix} 1 & 3 & 4 & 7 & 8 & 10 \\ 2 & 5 & 6 & 9 & & \end{pmatrix},$$

$$w_8 = w \begin{pmatrix} 1 & 2 & 4 & 6 & 8 & 10 \\ 3 & 5 & 7 & 9 & & \end{pmatrix},$$

$$w_9 = w \begin{pmatrix} 1 & 3 & 4 & 5 & 8 & 9 \\ 2 & 6 & 7 & 10 & & \end{pmatrix},$$

$$w_{10} = w \begin{pmatrix} 1 & 3 & 4 & 7 & 9 & 10 \\ 2 & 5 & 6 & 8 & & \end{pmatrix};$$

$\lambda = (5, 5)$:

$$w_1 = w \begin{pmatrix} 1 & 3 & 5 & 7 & 9 \\ 2 & 4 & 6 & 8 & 10 \end{pmatrix},$$

$$w_2 = w \begin{pmatrix} 1 & 3 & 5 & 7 & 8 \\ 2 & 4 & 6 & 9 & 10 \end{pmatrix},$$

$$w_3 = w \begin{pmatrix} 1 & 3 & 5 & 6 & 7 \\ 2 & 4 & 8 & 9 & 10 \end{pmatrix},$$

$$w_4 = w \begin{pmatrix} 1 & 2 & 3 & 4 & 9 \\ 5 & 6 & 7 & 8 & 10 \end{pmatrix}.$$

Proof. For each $\lambda \vdash n$, the number of highest weight vectors w_i given above, coincides with the multiplicity of $W(\lambda)$ in U_n . The explicit form of w_i is obtained by direct calculations. It is sufficient to prove that the w_i s are linearly independent in U_n . For most of the cases it is almost obvious. In the more complicated cases the linear independence is obtained by calculating the rank of the matrix with entries equal to the coefficients of some of $\text{tr}(x^{a_1}y^{b_1} \dots x^{a_p}y^{b_p})$ in the expression of w_i . For example, for $\lambda = (6, 3)$ we compare the coefficients of

$$\text{tr}(x^6y^3), \text{tr}(x^5y^2xy), \text{tr}(x^5yxy^2), \text{tr}(x^4y^2x^2y), \text{tr}(x^4yx^2y^2), \text{tr}(x^3yx^2yxy)$$

in the presentation of $w_1, \dots, w_6$ and obtain the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & -1 & -1 & -1 & 2 \\ -1 & -1 & -1 & 0 & -1 & -2 \\ -1 & 0 & 1 & 0 & 1 & 0 \\ 1 & 1 & -1 & 2 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

Its rank is equal to 6, and this gives that the trace polynomials $w_1, \dots, w_6$ are linearly independent in U_9 . $\square$

3. THE NEW GENERATORS

We fix the generic 4×4 matrices $X = (x_{ij})$ and $Y = (y_{ij})$. Changing the variables on the diagonals of X and Y , we can present them in the form

$$(5) \quad X = \frac{1}{4}\text{tr}(X)e + x, \quad Y = \frac{1}{4}\text{tr}(Y)e + y,$$

where e is the identity 4×4 matrix and x, y are generic traceless matrices. As in the case of generic matrices without restrictions on the trace, we may assume that x is diagonal. We fix the notation

$$x = \begin{pmatrix} x_1 & 0 & 0 & 0 \\ 0 & x_2 & 0 & 0 \\ 0 & 0 & x_3 & 0 \\ 0 & 0 & 0 & -(x_1 + x_2 + x_3) \end{pmatrix},$$

$$y = \begin{pmatrix} y_{11} & y_{12} & y_{13} & y_{14} \\ y_{21} & y_{22} & y_{23} & y_{24} \\ y_{31} & y_{32} & y_{33} & y_{34} \\ y_{41} & y_{42} & y_{43} & -(y_{11} + y_{22} + y_{33}) \end{pmatrix},$$

where x_1, x_2, x_3, y_{ij} are algebraically independent commuting variables. The algebra C_{42} is generated by $\text{tr}(X), \text{tr}(Y)$ and the traces of products of x and y . We denote by C_0 the algebra generated by $\text{tr}(z_1 \cdots z_k)$, $z_i = x, y$, and $k \geq 2$. Clearly,

$$(6) \quad C_{42} \cong K[\text{tr}(X), \text{tr}(Y)] \otimes_K C_0,$$

and it is sufficient to find a minimal system of generators of C_0 . Our idea is the following. We want to build inductively the GL_2 -module G of the a minimal system of generators. We start with the GL_2 -module $G_1 = W(1)$ generated by $\text{tr}(X)$, and further, we assume that the generators of higher degree are in C_0 . Assume that we have already found the module $G_2 \oplus \cdots \oplus G_{n-1}$ of generators of degree $< n$ of C_0 . Then we consider the symmetric algebra $K[G_2 \oplus \cdots \oplus G_{n-1}]$ and all trace identities for $M_4(K)$ of the form $f(x, y) - g(x, y) = 0$, where $f(x, y) \in K[G_2 \oplus \cdots \oplus G_{n-1}]$ and $g(x, y) \in U_n$. The elements $g(x, y)$ generate a GL_2 -submodule R_n of U_n . Then the GL_2 -submodule G_n of U_n is a complement of R_n in U_n . We may identify G_n with a GL_2 -submodule of C_0 . We add G_n to the module of generators and call it the GL_2 -module of new generators of degree n . Then $G_2 \oplus \cdots \oplus G_{10}$ is a GL_2 -module of generators of the algebra of invariants of two traceless 4×4 matrices and

$G = G_1 \oplus (G_2 \oplus \cdots \oplus G_{10})$ is a minimal GL_2 -module of generators of C_{42} . Let the homogeneous component of degree n of $K[G_2 \oplus \cdots \oplus G_{n-1}]$ be

$$K[G_2 \oplus \cdots \oplus G_{n-1}]^{(n)} = \sum_{(\lambda_1, \lambda_2) \vdash n} p(\lambda) W(\lambda).$$

Similarly, we have the decomposition

$$U_n = \sum_{(\lambda_1, \lambda_2) \vdash n} q(\lambda) W(\lambda)$$

from Proposition 2.1 and the decomposition

$$R_n = \sum_{(\lambda_1, \lambda_2) \vdash n} r(\lambda) W(\lambda) \subset U_n,$$

with unknown multiplicities $r(\lambda)$. If $v_1, \dots, v_{p(\lambda)} \in K[G_2 \oplus \cdots \oplus G_{n-1}]^{(n)}$ are linearly independent highest weight vectors of the direct sum of $p = p(\lambda)$ isomorphic copies of $W(\lambda)$ and $w_1, \dots, w_q \in U_n$, $q = q(\lambda)$, are linearly independent highest weight vectors in $qW(\lambda) \in U_n$, then the highest weight vectors of the GL_2 -submodule $rW(\lambda) \subset R_n$, $r = r(\lambda)$, are all trace identities of the form

$$w(x, y) = (\xi_1 v_1 + \cdots + \xi_p v_p) - (\eta_1 w_1 + \cdots + \eta_q w_q) = 0,$$

where $\xi_1, \dots, \xi_p, \eta_1, \dots, \eta_q$ are constants from the field K . We consider the elements $\xi_1, \dots, \xi_p, \eta_1, \dots, \eta_q$ as unknowns. The entries of the evaluation of $w(x, y)$ on the generic traceless matrices x, y are equal to 0. In this way we obtain a linear homogeneous system with respect to the unknowns $\xi_1, \dots, \xi_p, \eta_1, \dots, \eta_q$, and with coefficients which are polynomials in x_1, x_2, x_3 and y_{ij} . The module R_n is generated by all $\eta_1 w_1 + \cdots + \eta_q w_q$ which correspond to solutions $\xi_1, \dots, \xi_p, \eta_1, \dots, \eta_q$.

The highest weight vectors $w_1, \dots, w_q$ are given in the previous section. In order to find $v_1, \dots, v_p$, we need first to know the multiplicity $p(\lambda)$ of $W(\lambda)$ in $K[G_2 \oplus \cdots \oplus G_{n-1}]^{(n)}$. If $G_2 \oplus \cdots \oplus G_{n-1} = W_1 \oplus \cdots \oplus W_m$, where the GL_2 -modules $W_1, \dots, W_m$ are irreducible, then

$$K[G_2 \oplus \cdots \oplus G_{n-1}] = K[W_1] \otimes \cdots \otimes K[W_m].$$

If W_i is homogeneous of degree d_i , and $W_i^{\otimes_s b_i}$ denotes the b_i symmetric power of W_i , then

$$K[G_2 \oplus \cdots \oplus G_{n-1}]^{(n)} = \bigoplus \left(W_1^{\otimes_s b_1} \otimes \cdots \otimes W_m^{\otimes_s b_m} \right),$$

where the direct sum is on all $(b_1, \dots, b_m)$ with $b_1 d_1 + \cdots + b_m d_m = n$. The decomposition of $W_i^{\otimes_s b_i}$ can be obtained from the Hilbert series of $K[W_i]$ and this of the usual tensor product by the Littlewood-Richardson rule. From computational point of view it is convenient to calculate the homogeneous component of degree n of the Hilbert series of $K[G_2 \oplus \cdots \oplus G_{n-1}]$. If

$$H(W_i, t, u) = a_{i0} t^{d_i} + a_{i1} t^{d_i-1} u + \cdots + a_{i, d_i-1} t u^{d_i-1} + a_{i d_i} u^{d_i}, \quad a_{ij} \geq 0,$$

is the Hilbert series of W_i , then

$$H(K[G_2 \oplus \cdots \oplus G_{n-1}], t, u) = \prod_{i=1}^m \prod_{j=0}^{d_i} \frac{1}{(1 - t^j u^{d_i-j})^{a_{ij}}}.$$

Then the homogeneous component of degree n of the Hilbert series can be obtained using (3). The highest weight vectors can be found using the constructive approach with the derivations. We start the realization of this scheme step by step.

Lemma 3.1. *For $n \leq 5$, the GL_2 -modules G_n of the new generators of C_0 are*

$$G_2 = W(2, 0), \quad G_3 = W(3, 0), \quad G_4 = W(4, 0) \oplus W(2, 2), \quad G_5 = W(3, 2).$$

Proof. By Proposition 2.1

$$U_2 \oplus U_3 \oplus U_4 \oplus U_5 = W(2) \oplus W(3) \oplus W(4) \oplus W(2, 2) \oplus W(5, 0) \oplus W(3, 2).$$

Hence $G_2 \oplus G_3 \oplus G_4 \oplus G_5$ is a homomorphic image of this module. We shall show that $W(5, 0)$ belongs to the subalgebra generated by $G_2 \oplus G_3 \oplus G_4$. Since $W(5, 0)$ is generated by $\text{tr}(x^5)$, it is sufficient to show that $\text{tr}(x^5)$ can be expressed in C_0 in terms of $\text{tr}(x^2)$, $\text{tr}(x^3)$, and $\text{tr}(x^4)$. By the Cayley-Hamilton theorem, x satisfies an equation of degree 4 with coefficients depending on $\text{tr}(x^2)$, $\text{tr}(x^3)$, and $\text{tr}(x^4)$. Direct verification gives that

$$x^4 - \frac{1}{2}\text{tr}(x^2)x^2 - \frac{1}{3}\text{tr}(x^3)x + \left(-\frac{1}{4}\text{tr}^2(x^2) + \frac{1}{8}\text{tr}(x^4)\right)e = 0.$$

Multiplying by x , taking the trace, and using that $\text{tr}(x) = 0$, we obtain

$$(7) \quad \text{tr}(x^5) - \frac{5}{6}\text{tr}(x^2)\text{tr}(x^3) = 0.$$

(In the same way, by easy induction, we obtain that $\text{tr}(x^n) \in K[\text{tr}(x^2), \text{tr}(x^3), \text{tr}(x^4)]$ in C_0 for any $n \geq 5$.) Hence, the GL_2 -module of the new generators of C_0 , of degree ≤ 5 , is a homomorphic image of the module

$$F = W(2) \oplus W(3) \oplus W(4) \oplus W(2, 2) \oplus W(3, 2).$$

The Hilbert series of F is

$$\begin{aligned} H(F, t, u) &= (t^2 + tu + u^2) + (t^3 + t^2u + tu^2 + u^3) \\ &\quad + (t^4 + t^3u + 2t^2u^2 + tu^3 + u^4) + (t^3u^2 + t^2u^3) \end{aligned}$$

and the Hilbert series of the symmetric algebra of F is

$$H(K[F], t, u) = \frac{1}{q_2(t, u)q_3(t, u)q_4(t, u)q_5(t, u)},$$

$$(8) \quad \begin{aligned} q_2(t, u) &= (1 - t^2)(1 - tu)(1 - u^2), \\ q_3(t, u) &= (1 - t^3)(1 - t^2u)(1 - tu^2)(1 - u^3), \\ q_4(t, u) &= (1 - t^4)(1 - t^3u)(1 - t^2u^2)^2(1 - tu^3)(1 - u^4), \\ q_5(t, u) &= (1 - t^3u^2)(1 - t^2u^3). \end{aligned}$$

The algebra $K[F]$ maps naturally on the subalgebra of C_0 generated by $G_2 \oplus G_3 \oplus G_4 \oplus G_5$, and the coefficients of the Hilbert series of $K[F]$ are bigger or equal to the corresponding coefficients of the Hilbert series of this subalgebra. By direct calculation, we compare the coefficients of the homogeneous components of degree ≤ 5 of the Hilbert series of $H(K[F], t, u)$ and $H(C_0, t, u)$. We find that they coincide and this implies that the homogeneous components of degree ≤ 5 of $K[F]$ and C_0 are isomorphic as GL_2 -modules, completing the proof of the lemma. $\square$

In the following lemmas the elements w_i are taken from Lemma 2.2. The relations have been found using the scheme described in the beginning of the section. The proofs are direct verifications, performed by Maple.

Lemma 3.2. *For $\lambda = (4, 2)$, the following is a relation in C_0 :*

$$6w_1 - 12w_2 + 6v_1 + 2v_2 - 3v_3 - 5v_4 = 0,$$

where

$$\begin{aligned} v_1 &= \text{tr}(x^3)\text{tr}(xy^2) - \text{tr}^2(x^2y), \\ v_2 &= 3\text{tr}(x^4)\text{tr}(y^2) - 6\text{tr}(x^3y)\text{tr}(xy) + (2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^2), \\ v_3 &= \text{tr}(x^2)(\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy)), \\ v_4 &= \frac{1}{2}\text{tr}([x, y]^2)\text{tr}(x^2). \end{aligned}$$

Lemma 3.3. *The following relations hold in C_0 :*

For $\lambda = (5, 2)$:

$$\begin{aligned} -24w_1 + 18v_1 + 4v_2 + 16v_3 - 3v_4 - 4v_5 &= 0, \\ 24w_2 + 6v_1 - 4v_2 + 3v_4 + 4v_5 &= 0, \end{aligned}$$

where

$$\begin{aligned} v_1 &= \text{tr}([x, y]^2x)\text{tr}(x^2), \\ v_2 &= 3\text{tr}(x^4)\text{tr}(xy^2) - 6\text{tr}(x^3y)\text{tr}(x^2y) + (\text{tr}(xyxy) + 2\text{tr}(x^2y^2))\text{tr}(x^3), \\ v_3 &= \frac{1}{2}(\text{tr}([x, y]^2)\text{tr}(x^3), \\ v_4 &= \text{tr}(x^2)(\text{tr}(x^2)\text{tr}(xy^2) - 2\text{tr}(xy)\text{tr}(x^2y) + \text{tr}(y^2)\text{tr}(x^3)), \\ v_5 &= (-\text{tr}(x^2))\text{tr}(y^2) + \text{tr}^2(xy)\text{tr}(x^3). \end{aligned}$$

For $\lambda = (4, 3)$:

$$4w_1 - 8w_2 - 10v_1 - 4v_2 + 3v_3 = 0,$$

where

$$\begin{aligned} v_1 &= \text{tr}([x, y]^2x)\text{tr}(xy) - \text{tr}([x, y]^2y)\text{tr}(x^2), \\ v_2 &= \text{tr}(x^4)\text{tr}(y^3) - 3\text{tr}(x^3y)\text{tr}(xy^2) \\ &\quad + (\text{tr}(xyxy) + 2\text{tr}(x^2y^2))\text{tr}(x^2y) - \text{tr}(xy^3)\text{tr}(x^3), \\ v_3 &= \text{tr}^2(x^2)\text{tr}(y^3) - 3\text{tr}(x^2)\text{tr}(xy)\text{tr}(xy^2) \\ &\quad + \text{tr}(x^2)\text{tr}(y^2)\text{tr}(x^2y) + 2\text{tr}^2(xy)\text{tr}(x^2y) - \text{tr}(xy)\text{tr}(y^2)\text{tr}(x^3). \end{aligned}$$

Lemma 3.4. *The following relations hold in C_0 :*

For $\lambda = (6, 2)$:

$$\begin{aligned} 12w_1 - 6v_1 - 4v_2 - 6v_4 + 3v_7 &= 0, \\ 24w_2 - 6v_1 + 8v_2 - 6v_5 - 2v_6 + 5v_7 - 6v_8 + 6v_9 + 3v_{10} &= 0, \\ 48w_3 + 18v_1 - 40v_2 - 16v_3 - 8v_4 + 18v_5 + 10v_6 - 13v_7 + 2v_8 - 6v_9 - 9v_{10} &= 0, \end{aligned}$$

where

$$\begin{aligned} v_1 &= \text{tr}([x, y]^2x^2)\text{tr}(x^2), \\ v_2 &= \text{tr}([x, y]^2x)\text{tr}(x^3), \\ v_3 &= \text{tr}(x^4)(2\text{tr}(x^2y^2) + \text{tr}(xyxy)) - 3\text{tr}^2(x^3y), \\ v_4 &= \frac{1}{2}\text{tr}([x, y]^2)\text{tr}(x^4), \\ v_5 &= \text{tr}(x^4)(\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy)), \\ v_6 &= 3\text{tr}(x^4)\text{tr}^2(xy) - 6\text{tr}(x^3y)\text{tr}(x^2)\text{tr}(xy) + (2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}^2(x^2), \\ v_7 &= \frac{1}{2}\text{tr}([x, y]^2)\text{tr}^2(x^2), \\ v_8 &= \text{tr}(x^3)(\text{tr}(x^3)\text{tr}(y^2) - 2\text{tr}(x^2y)\text{tr}(xy) + \text{tr}(xy^2)\text{tr}(x^2)), \end{aligned}$$

$$\begin{aligned} v_9 &= \text{tr}^2(x^3)\text{tr}(y^2) - 2\text{tr}(x^3)\text{tr}(x^2y)\text{tr}(xy) + \text{tr}^2(x^2y)\text{tr}(x^2), \\ v_{10} &= (\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy))\text{tr}^2(x^2). \end{aligned}$$

For $\lambda = (5, 3)$:

$$\begin{aligned} 16w_1 - 8w_2 - v_1 - 4v_2 - 4v_3 + 2v_4 - v_5 &= 0, \\ 2w_3 - v_2 &= 0, \end{aligned}$$

where

$$\begin{aligned} v_1 &= -2\text{tr}([x, y]^2x^2)\text{tr}(xy) + (\text{tr}([x, y]^2xy) + \text{tr}([x, y]^2yx))\text{tr}(x^2), \\ v_2 &= (\text{tr}(x^2y^2xy) - \text{tr}(y^2x^2yx))\text{tr}(x^2), \\ v_3 &= \text{tr}([x, y]^2x)\text{tr}(x^2y) - \text{tr}([x, y]^2y)\text{tr}(x^3), \\ v_4 &= \text{tr}(x^4)\text{tr}(xy)\text{tr}(y^2) - \text{tr}(x^3y)\text{tr}(x^2)\text{tr}(y^2) \\ &\quad - 2\text{tr}(x^3y)\text{tr}^2(xy) + (2\text{tr}(x^2y^2) + \text{tr}(xyxy)\text{tr}(x^2)\text{tr}(xy) - \text{tr}(xy^3)\text{tr}^2(x^2)), \\ v_5 &= -2\text{tr}(x^3)\text{tr}(xy^2)\text{tr}(xy) + 2\text{tr}(x^2y)\text{tr}(x^2y)\text{tr}(xy) \\ &\quad + \text{tr}(x^3)\text{tr}(y^3)\text{tr}(x^2) - \text{tr}(x^2y)\text{tr}(xy^2)\text{tr}(x^2). \end{aligned}$$

For $\lambda = (4, 4)$:

$$\begin{aligned} -24w_1 + 12w_2 - 12v_1 - 4v_2 + 16v_3 + v_4 + 16v_5 &= 0, \\ -72w_1 + 24w_3 - 42v_1 - 20v_2 + 56v_3 + 3v_4 + 66v_5 + 6v_6 + 6v_7 &= 0, \end{aligned}$$

where

$$\begin{aligned} v_1 &= \text{tr}([x, y]^2x^2)\text{tr}(y^2) - (\text{tr}([x, y]^2xy) + \text{tr}([x, y]^2yx))\text{tr}(xy) + \text{tr}([x, y]^2y^2)\text{tr}(x^2), \\ v_2 &= 3\text{tr}(x^4)\text{tr}(y^4) - 12\text{tr}(x^3y)\text{tr}(xy^3) + (2\text{tr}(x^2y^2) + \text{tr}(xyxy))^2, \\ v_3 &= \frac{1}{4}\text{tr}^2([x, y]^2), \\ v_4 &= 3\text{tr}(x^4)\text{tr}^2(y^2) - 12\text{tr}(x^3y)\text{tr}(xy)\text{tr}(y^2) \\ &\quad + 2(2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^2)\text{tr}(y^2) + 4(2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}^2(xy) \\ &\quad - 12\text{tr}(xy^3)\text{tr}(x^2)\text{tr}(xy) + 3\text{tr}(y^4)\text{tr}^2(x^2), \\ v_5 &= \frac{1}{2}\text{tr}([x, y]^2)(\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy)), \\ v_6 &= -\text{tr}(x^3)\text{tr}(xy^2)\text{tr}(y^2) + \text{tr}^2(x^2y)\text{tr}(y^2) + \text{tr}(x^3)\text{tr}(y^3)\text{tr}(xy) \\ &\quad - \text{tr}(x^2y)\text{tr}(xy^2)\text{tr}(xy) - \text{tr}(x^2y)\text{tr}(y^3)\text{tr}(x^2) + \text{tr}^2(xy^2)\text{tr}(x^2), \\ v_7 &= (\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy))^2. \end{aligned}$$

Lemma 3.5. *The following relations hold in C_0 between the elements w_i and the elements v_j given below:*

For $\lambda = (7, 2)$:

$$\begin{aligned} 48w_1 - 16v_1 - 12v_2 - 12v_3 - 4v_5 - 16v_7 - v_9 + 4v_{10} &= 0, \\ 144w_2 - 24v_1 + 36v_2 - 24v_4 - 20v_5 + 48v_6 + 20v_7 - 24v_8 + 9v_9 &= 0, \\ 48w_3 - 12v_2 - 4v_5 - 8v_7 - v_9 + 4v_{10} &= 0, \\ v_1 &= \text{tr}([x, y]^2x^2)\text{tr}(x^3), \\ v_2 &= \text{tr}([x, y]^2x)\text{tr}(x^4), \\ v_3 &= \text{tr}([x, y]^2x)\text{tr}^2(x^2), \\ v_4 &= \text{tr}(x^4)(\text{tr}(x^3)\text{tr}(y^2) + \text{tr}(xy^2)\text{tr}(x^2) - 2\text{tr}(x^2y)\text{tr}(xy)), \\ v_5 &= 3\text{tr}(x^4)\text{tr}(xy^2)\text{tr}(x^2) \end{aligned}$$

$$\begin{aligned}
& +(2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^3)\text{tr}(x^2) - 6\text{tr}(x^3y)\text{tr}(x^2y)\text{tr}(x^2), \\
v_6 &= \text{tr}(x^4)\text{tr}(xy^2)\text{tr}(x^2) - \text{tr}(x^4)\text{tr}(x^2y)\text{tr}(xy) \\
& + \text{tr}(x^3y)\text{tr}(x^3)\text{tr}(xy) - \text{tr}(x^3y)\text{tr}(x^2y)\text{tr}(x^2), \\
v_7 &= \frac{1}{2}\text{tr}([x, y]^2)\text{tr}(x^3)\text{tr}(x^2), \\
v_8 &= \text{tr}(x^3)(\text{tr}(x^3)\text{tr}(xy^2) - \text{tr}^2(x^2y)), \\
v_9 &= (\text{tr}(x^3)\text{tr}(y^2) - 2\text{tr}(x^2y)\text{tr}(xy) + \text{tr}(xy^2)\text{tr}(x^2))\text{tr}^2(x^2), \\
v_{10} &= (\text{tr}(x^3)\text{tr}^2(xy) - 2\text{tr}(x^2y)\text{tr}(x^2)\text{tr}(xy) + \text{tr}(xy^2)\text{tr}^2(x^2))\text{tr}(x^2).
\end{aligned}$$

For $\lambda = (6, 3)$:

$$\begin{aligned}
& -24w_1 + 48w_2 - 12v_1 + 3v_2 - 24v_3 + 36v_4 + 12v_5 \\
& -6v_6 - 2v_7 + 8v_8 + 7v_9 + 3v_{10} - 15v_{11} - 7v_{12} = 0, \\
& 2w_3 + v_1 = 0, \\
24w_4 - 12v_1 - 12v_2 - 12v_3 - 24v_4 + 6v_5 - 4v_7 + 4v_8 - 16v_9 - 6v_{11} - v_{12} &= 0, \\
w_1 + w_5 - v_1 - v_3 &= 0, \\
24w_1 + 48w_6 + 3v_2 - 24v_3 - 12v_4 - 18v_5 \\
-6v_6 - 10v_7 - 4v_8 - 9v_9 + 3v_{10} + 3v_{11} + 6v_{12} &= 0, \\
v_1 &= \text{tr}([x, y]^3x)\text{tr}(x^2), \\
v_2 &= 2\text{tr}([x, y]x^2)\text{tr}(x^2y) - (\text{tr}([x, y]^2xy) + \text{tr}([x, y]^2yx))\text{tr}(x^3), \\
v_3 &= (\text{tr}(x^2y^2xy) - \text{tr}(y^2x^2yx))(\text{tr}(x^3)), \\
v_4 &= \text{tr}([x, y]^2x)\text{tr}(x^3y) - \text{tr}([x, y]^2y)\text{tr}(x^4), \\
v_5 &= (\text{tr}([x, y]^2x)\text{tr}(xy) - \text{tr}([x, y]^2y)\text{tr}(x^2))\text{tr}(x^2), \\
v_6 &= (\text{tr}(x^4)\text{tr}(x^2y) - \text{tr}(x^3y)\text{tr}(x^3))\text{tr}(y^2) \\
& + 2(-\text{tr}(x^4)\text{tr}(xy^2) + \text{tr}(x^3y)\text{tr}(x^2y))\text{tr}(xy) \\
& + (2\text{tr}(x^3y)\text{tr}(xy^2) - (2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^2y) + \text{tr}(xy^3)\text{tr}(x^3))\text{tr}(x^2), \\
v_7 &= (3\text{tr}(x^4)\text{tr}(xy^2) - 6\text{tr}(x^3y)\text{tr}(x^2y) + (2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^3))\text{tr}(xy) \\
& + (-3\text{tr}(x^3y)\text{tr}(xy^2) + 2(2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^2y) - 3\text{tr}(xy^3)\text{tr}(x^3))\text{tr}(x^2), \\
v_8 &= (\text{tr}(x^4)\text{tr}(y^3) - 3\text{tr}(x^3y)\text{tr}(xy^2) \\
& + (2\text{tr}(x^2y^2) + \text{tr}(xyxy))\text{tr}(x^2y) - \text{tr}(xy^3)\text{tr}(x^3))\text{tr}(x^2), \\
v_9 &= \frac{1}{2}\text{tr}([x, y]^2)(\text{tr}(x^3)\text{tr}(xy) - \text{tr}(x^2y)\text{tr}(x^2)), \\
v_{10} &= \text{tr}^2(x^3)\text{tr}(y^3) - 3\text{tr}(x^3)\text{tr}(x^2y)\text{tr}(xy^2) + 2\text{tr}^3(x^2y), \\
v_{11} &= \text{tr}(x^3)\text{tr}(x^2)\text{tr}(xy)\text{tr}(y^2) - \text{tr}(x^3)\text{tr}^3(xy) \\
& - \text{tr}(x^2y)\text{tr}^2(x^2)\text{tr}(y^2) + \text{tr}(x^2y)\text{tr}(x^2)\text{tr}^2(xy), \\
v_{12} &= -3\text{tr}(x^3)\text{tr}(x^2)\text{tr}(xy)\text{tr}(y^2) + 2\text{tr}(x^3)\text{tr}^3(xy) \\
& + 3\text{tr}(x^2y)\text{tr}^2(x^2)\text{tr}(y^2) - 3\text{tr}(xy^2)\text{tr}^2(x^2)\text{tr}(xy) + \text{tr}(y^3)\text{tr}^3(x^2).
\end{aligned}$$

For $\lambda = (5, 4)$:

$$\begin{aligned}
& w_1 - v_3 = 0, \\
24w_2 + 12v_1 - 6v_2 + 12v_3 - 6v_4 - 2v_5 + 2v_6 + v_7 &= 0, \\
w_3 + v_1 &= 0, \\
24w_4 + 6v_2 + 12v_4 - 4v_6 + v_7 + 3v_8 &= 0, \\
v_1 &= \text{tr}([x, y]^3x)\text{tr}(xy) - \text{tr}([x, y]^3y)\text{tr}(x^2), \\
v_2 &= \text{tr}([x, y]^2x^2)\text{tr}(xy^2)
\end{aligned}$$

$$\begin{aligned}
& -(\operatorname{tr}([x, y]^2 xy) + \operatorname{tr}([x, y]^2 yx))\operatorname{tr}(x^2 y) + \operatorname{tr}([x, y]^2 y^2)\operatorname{tr}(x^3), \\
v_3 &= \frac{1}{2}\operatorname{tr}([x, y]^2)\operatorname{tr}([x, y]^2 x), \\
v_4 &= \operatorname{tr}([x, y]^2 x)(\operatorname{tr}(x^2)\operatorname{tr}(y^2) - \operatorname{tr}^2(xy)), \\
v_5 &= (3\operatorname{tr}(x^4)\operatorname{tr}(xy^2) - 6\operatorname{tr}(x^3 y)\operatorname{tr}(x^2 y) + (2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}(x^3))\operatorname{tr}(y^2) \\
& \quad + (-\operatorname{tr}(x^4)\operatorname{tr}(y^3) - 3\operatorname{tr}(x^3 y)\operatorname{tr}(xy^2) \\
& \quad + 3(2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}(x^2 y) - 5\operatorname{tr}(xy^3)\operatorname{tr}(x^3))\operatorname{tr}(xy) \\
& \quad + (\operatorname{tr}(x^3 y)\operatorname{tr}(y^3) - 3\operatorname{tr}(xy^3)\operatorname{tr}(x^2 y) + 2\operatorname{tr}(y^4)\operatorname{tr}(x^3))\operatorname{tr}(x^2), \\
v_6 &= (\operatorname{tr}(x^4)\operatorname{tr}(y^3) - 3\operatorname{tr}(x^3 y)\operatorname{tr}(xy^2) \\
& \quad + (2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}(x^2 y) - \operatorname{tr}(xy^3)\operatorname{tr}(x^3))\operatorname{tr}(xy) \\
& \quad + (-\operatorname{tr}(x^3 y)\operatorname{tr}(y^3) + (2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}(xy^2) \\
& \quad - 3\operatorname{tr}(xy^3)\operatorname{tr}(x^2 y) + \operatorname{tr}(y^4)\operatorname{tr}(x^3))\operatorname{tr}(x^2), \\
v_7 &= \frac{1}{2}\operatorname{tr}([x, y]^2)(\operatorname{tr}(x^3)\operatorname{tr}(y^2) - 2\operatorname{tr}(x^2 y)\operatorname{tr}(xy) + \operatorname{tr}(xy^2)\operatorname{tr}(x^2)), \\
v_8 &= (\operatorname{tr}(x^2)\operatorname{tr}(y^2) - \operatorname{tr}^2(xy))(\operatorname{tr}(x^3)\operatorname{tr}(y^2) - 2\operatorname{tr}(x^2 y)\operatorname{tr}(xy) + \operatorname{tr}(xy^2)\operatorname{tr}(x^2)).
\end{aligned}$$

Lemma 3.6. *For $\lambda = (8, 2)$ the elements w_i and v_j satisfy the following relations in C_0 :*

$$\begin{aligned}
& 360w_1 - 90v_1 - 45v_2 - 150v_3 - 90v_6 - 4v_7 - 36v_8 \\
& -80v_9 - 45v_{12} + v_{13} - 10v_{14} + 9v_{15} = 0, \\
& 720w_2 + 90v_1 + 90v_2 - 420v_3 + 90v_4 - 90v_5 - 135v_6 + 90v_8 \\
& -80v_9 + 12v_{10} + 9v_{11} + 60v_{12} - 2v_{13} - 20v_{14} - 18v_{15} - 45v_{16} = 0, \\
& 720w_3 - 90v_1 - 180v_2 + 90v_4 + 150v_5 - 195v_6 + 8v_7 \\
& + 162v_8 - 12v_{10} - 99v_{11} - 150v_{12} + 40v_{14} + 45v_{16} = 0, \\
& 1440w_4 + 270v_2 - 960v_3 - 240v_5 - 120v_6 - 16v_7 - 144v_8 \\
& -320v_9 + 30v_{10} + 180v_{11} + 195v_{12} - 80v_{14} + 90v_{15} - 135v_{16} = 0,
\end{aligned}$$

where

$$\begin{aligned}
v_1 &= \operatorname{tr}([x, y]^2 x^2)\operatorname{tr}(x^4), \\
v_2 &= \operatorname{tr}([x, y]^2 x^2)\operatorname{tr}^2(x^2), \\
v_3 &= \operatorname{tr}([x, y]^2 x)\operatorname{tr}(x^3)\operatorname{tr}(x^2), \\
v_4 &= \operatorname{tr}^2(x^3 y)\operatorname{tr}(x^2) - 2\operatorname{tr}(x^4)\operatorname{tr}(x^3 y)\operatorname{tr}(xy) + \operatorname{tr}^2(x^4)(y^2), \\
v_5 &= (2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}(x^4)\operatorname{tr}(x^2) - 3\operatorname{tr}^2(x^3 y)\operatorname{tr}(x^2), \\
v_6 &= \frac{1}{2}\operatorname{tr}([x, y]^2)\operatorname{tr}(x^4)\operatorname{tr}(x^2), \\
v_7 &= 5(2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}^2(x^3) - 30\operatorname{tr}(x^3 y)\operatorname{tr}(x^2 y)\operatorname{tr}(x^3) \\
& \quad + 3\operatorname{tr}(x^4)(2\operatorname{tr}(xy^2)\operatorname{tr}(x^3) + 3\operatorname{tr}^2(x^2 y)), \\
v_8 &= \operatorname{tr}(x^4)(\operatorname{tr}(xy^2)\operatorname{tr}(x^3) - \operatorname{tr}^2(x^2 y)), \\
v_9 &= \frac{1}{2}\operatorname{tr}([x, y]^2)\operatorname{tr}^2(x^3), \\
v_{10} &= (5(2\operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy))\operatorname{tr}^2(x^2) - 30\operatorname{tr}(x^3 y)\operatorname{tr}(xy)\operatorname{tr}(x^2) \\
& \quad + 3\operatorname{tr}(x^4)(4\operatorname{tr}^2(xy) + \operatorname{tr}(y^2)\operatorname{tr}(x^2)))\operatorname{tr}(x^2), \\
v_{11} &= \operatorname{tr}(x^4)(\operatorname{tr}(y^2)\operatorname{tr}(x^2) - \operatorname{tr}^2(xy))\operatorname{tr}(x^2), \\
v_{12} &= (\operatorname{tr}(x^2 y^2) - \operatorname{tr}(xyxy))\operatorname{tr}^3(x^2), \\
v_{13} &= 5\operatorname{tr}^2(x^3)\operatorname{tr}(x^3)(\operatorname{tr}(y^2)\operatorname{tr}(x^2) + 2\operatorname{tr}^2(xy)) - 30\operatorname{tr}(x^3)\operatorname{tr}(x^2 y)\operatorname{tr}(xy)\operatorname{tr}(x^2)
\end{aligned}$$

$$\begin{aligned}
& +3(2\text{tr}(x^3)\text{tr}(xy^2) + 3\text{tr}^2(x^2y))\text{tr}^2(x^2), \\
v_{14} &= \text{tr}^2(x^3)(\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy)), \\
v_{15} &= (\text{tr}(x^3)\text{tr}(xy^2) - \text{tr}^2(x^2y))\text{tr}^2(x^2), \\
v_{16} &= (\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy))\text{tr}^3(x^2).
\end{aligned}$$

Lemma 3.7. *In C_0 and for $\lambda = (7, 3)$, the elements $w_1, \dots, w_7$ and*

$$\begin{aligned}
v_1 &= \text{tr}([x, y]^3 x^2) \text{tr}(x^2), \\
v_2 &= \text{tr}([x, y]^3 x) \text{tr}(x^3), \\
v_3 &= (\text{tr}([x, y]^2 xy) + \text{tr}([x, y]^2 yx)) \text{tr}(x^4) - 2\text{tr}([x, y]^2 x^2) \text{tr}(x^3 y), \\
v_4 &= (\text{tr}(x^2 y^2 xy) - \text{tr}(y^2 x^2 yx)) \text{tr}(x^4), \\
v_5 &= ((\text{tr}([x, y]^2 xy) + \text{tr}([x, y]^2 yx)) \text{tr}(x^2) - 2\text{tr}([x, y]^2 x^2) \text{tr}(xy)) \text{tr}(x^2), \\
v_6 &= (\text{tr}(x^2 y^2 xy) - \text{tr}(y^2 x^2 yx)) \text{tr}^2(x^2), \\
v_7 &= (\text{tr}([x, y]^2 x) \text{tr}(x^2 y) - \text{tr}([x, y]^2 y) \text{tr}(x^3)) \text{tr}(x^2), \\
v_8 &= \text{tr}([x, y]^2 x) (\text{tr}(x^3) \text{tr}(xy) - \text{tr}(x^2 y) \text{tr}(x^2)), \\
v_9 &= (-6\text{tr}^2(x^3 y) + 2\text{tr}(x^4) (2\text{tr}(x^2 y^2) + \text{tr}(xyxy))) \text{tr}(xy) \\
&+ (\text{tr}(x^3 y) (2\text{tr}(x^2 y^2) + \text{tr}(xyxy)) - 3\text{tr}(x^4) \text{tr}(xy^3)) \text{tr}(x^2), \\
v_{10} &= \frac{1}{2} \text{tr}([x, y]^2) (\text{tr}(x^4) \text{tr}(xy) - \text{tr}(x^3 y) \text{tr}(x^2)), \\
v_{11} &= \text{tr}(x^4) \text{tr}(x^3) \text{tr}(y^3) - 3\text{tr}(x^4) \text{tr}(x^2 y) \text{tr}(xy^2) + 6\text{tr}(x^3 y) \text{tr}^2(x^2 y) \\
&- 2(2\text{tr}(x^2 y^2) + \text{tr}(xyxy)) \text{tr}(x^3) \text{tr}(x^2 y) + 2\text{tr}(xy^3) \text{tr}^2(x^3), \\
v_{12} &= -\text{tr}(x^4) \text{tr}(x^2 y) \text{tr}(xy^2) + \text{tr}(x^3 y) \text{tr}(x^3) \text{tr}(xy^2) + 2\text{tr}(x^3 y) \text{tr}^2(x^2 y) \\
&- (2\text{tr}(x^2 y^2) + \text{tr}(xyxy)) \text{tr}(x^3) \text{tr}(x^2 y) + \text{tr}(xy^3) \text{tr}^2(x^3), \\
v_{13} &= -\text{tr}(x^4) \text{tr}(x^2) \text{tr}(xy) \text{tr}(y^2) + \text{tr}(x^4) \text{tr}^3(xy) \\
&+ \text{tr}(x^3 y) \text{tr}^2(x^2) \text{tr}(y^2) - \text{tr}(x^3 y) \text{tr}(x^2) \text{tr}^2(xy), \\
v_{14} &= \text{tr}(x^4) \text{tr}^3(xy - 3\text{tr}(x^3 y) \text{tr}(x^2) \text{tr}^2(xy) \\
&+ (2\text{tr}(x^2 y^2) + \text{tr}(xyxy)) \text{tr}^2(x^2) \text{tr}(xy) - \text{tr}(xy^3) \text{tr}^3(x^2), \\
v_{15} &= \text{tr}^2(x^3) \text{tr}(xy) \text{tr}(y^2) - \text{tr}(x^3) \text{tr}(x^2 y) \text{tr}(x^2) \text{tr}(y^2) - 2\text{tr}(x^3) \text{tr}(x^2 y) \text{tr}^2(xy) \\
&+ \text{tr}(x^3) \text{tr}(xy^2) \text{tr}(x^2) \text{tr}(xy) + 2\text{tr}^2(x^2 y) \text{tr}(x^2) \text{tr}(xy) - \text{tr}(x^2 y) \text{tr}(xy^2) \text{tr}^2(x^2), \\
v_{16} &= -2\text{tr}(x^3) \text{tr}(xy^2) \text{tr}(x^2) \text{tr}(xy) + 2\text{tr}^2(x^2 y) \text{tr}(x^2) \text{tr}(xy) \\
&+ \text{tr}(x^3) \text{tr}(y^3) \text{tr}^2(x^2 - \text{tr}(x^2 y) \text{tr}(xy^2) \text{tr}^2(x^2)),
\end{aligned}$$

satisfy the relations

$$\begin{aligned}
24w_1 - 12v_1 - 8v_2 - 18v_4 + 9v_6 &= 0, \\
48w_2 - 24v_2 + 12v_3 - 24v_4 - 3v_5 - 8v_7 + 4v_8 - 12v_{10} + 4v_{11} - 2v_{15} - v_{16} &= 0, \\
48w_3 - 24v_1 - 8v_2 + 12v_3 - 12v_4 - 3v_5 + 6v_6 - 8v_7 \\
&+ 4v_8 - 12v_{10} + 4v_{11} - 2v_{15} - v_{16} = 0, \\
12w_4 - 4v_2 - 3v_6 &= 0, \\
48w_5 - 60v_1 + 8v_2 + 6v_3 - 24v_4 + 36v_6 + 10v_7 \\
&+ 4v_8 - 4v_9 - 10v_{10} + 10v_{11} - 12v_{12} + 6v_{13} - 2v_{15} - v_{16} = 0, \\
48w_6 - 48v_1 - 8v_2 - 36v_4 - 3v_5 + 30v_6 - 32v_7 \\
&- 20v_8 - 8v_{11} + 24v_{12} - 6v_{13} + 6v_{14} - 6v_{15} + 3v_{16} = 0, \\
12w_7 - 6v_1 + 4v_2 + 3v_6 &= 0.
\end{aligned}$$

Lemma 3.8. *The relation*

$$\sum_{i=1}^{10} \alpha_i w_i + \sum_{j=1}^{24} \beta_j v_j = 0,$$

holds in C_0 between w_i and v_j for $\lambda = (6, 4)$, where

$$\begin{aligned} v_1 &= (\operatorname{tr}([x, y]^3 xy) + \operatorname{tr}([x, y]^3 yx)) \operatorname{tr}(x^2) - 2 \operatorname{tr}([x, y]^3 x^2) \operatorname{tr}(xy), \\ v_2 &= \operatorname{tr}([x, y]^4) \operatorname{tr}(x^2), \\ v_3 &= \operatorname{tr}([x, y]^3 x) \operatorname{tr}(x^2 y) - \operatorname{tr}([x, y]^3 y) \operatorname{tr}(x^3), \\ v_4 &= \operatorname{tr}([x, y]^2 x^2) (2 \operatorname{tr}(x^2 y^2) + \operatorname{tr}(xyxy)) \\ &\quad - 3 (\operatorname{tr}([x, y]^2 xy) + \operatorname{tr}([x, y]^2 yx)) \operatorname{tr}(x^3 y) + 3 \operatorname{tr}([x, y]^2 y^2) \operatorname{tr}(x^4), \\ v_5 &= \frac{1}{2} \operatorname{tr}([x, y]^2 x^2) \operatorname{tr}([x, y]^2), \\ v_6 &= \operatorname{tr}([x, y]^2 x^2) \operatorname{tr}^2(xy) \\ &\quad - (\operatorname{tr}([x, y]^2 yx) + \operatorname{tr}([x, y]^2 xy)) \operatorname{tr}(x^2) \operatorname{tr}(xy) + \operatorname{tr}([x, y]^2 y^2) \operatorname{tr}^2(x^2), \\ v_7 &= \operatorname{tr}([x, y]^2 x^2) (\operatorname{tr}(x^2) \operatorname{tr}(y^2) - \operatorname{tr}^2(xy)), \\ v_8 &= \operatorname{tr}^2 t([x, y]^2 x), \\ v_9 &= \operatorname{tr}([x, y]^2 x) (\operatorname{tr}(x^3) \operatorname{tr}(y^2) - 2 \operatorname{tr}(x^2 y) \operatorname{tr}(xy) + \operatorname{tr}(xy^2) \operatorname{tr}(x^2)), \\ v_{10} &= \operatorname{tr}([x, y]^2 x) (-\operatorname{tr}(x^3) \operatorname{tr}(y^2) + \operatorname{tr}(x^2 y) \operatorname{tr}(xy)) \\ &\quad + \operatorname{tr}([x, y]^2 y) (\operatorname{tr}(x^3) \operatorname{tr}(xy) - \operatorname{tr}(x^2 y) \operatorname{tr}(x^2)), \\ v_{11} &= (\operatorname{tr}(x^4) \operatorname{tr}(x^2 y^2) + \operatorname{tr}(x^4) \operatorname{tr}(xyxy) - 2 \operatorname{tr}^2(x^3 y)) \operatorname{tr}(y^2) \\ &\quad + (-2 \operatorname{tr}(x^4) \operatorname{tr}(xy^3) + 2 \operatorname{tr}(x^3 y) \operatorname{tr}(x^2 y^2)) \operatorname{tr}(xy) \\ &\quad + (\operatorname{tr}(x^4) \operatorname{tr}(y^4) - 2 \operatorname{tr}(x^3 y) \operatorname{tr}(xy^3) + \operatorname{tr}(x^2 y^2) \operatorname{tr}(xyxy)) \operatorname{tr}(x^2), \\ v_{12} &= (\operatorname{tr}(x^4) \operatorname{tr}(x^2 y^2) - \operatorname{tr}^2(x^3 y)) \operatorname{tr}(y^2) \\ &\quad + (-\operatorname{tr}(x^4) \operatorname{tr}(xy^3) + \operatorname{tr}(x^3 y) \operatorname{tr}(xyxy)) \operatorname{tr}(xy) \\ &\quad + (\operatorname{tr}(x^3 y) \operatorname{tr}(xy^3) - \operatorname{tr}(x^2 y^2) \operatorname{tr}(xyxy)) \operatorname{tr}(x^2), \\ v_{13} &= (\operatorname{tr}(x^4) \operatorname{tr}(y^4) - 4 \operatorname{tr}(x^3 y) \operatorname{tr}(xy^3) + \operatorname{tr}^2(x^2 y^2) + 2 \operatorname{tr}(x^2 y^2) \operatorname{tr}(xyxy)) \operatorname{tr}(x^2), \\ v_{14} &= (-\operatorname{tr}(x^4) \operatorname{tr}(y^4) + 4 \operatorname{tr}(x^3 y) \operatorname{tr}(xy^3) - 4 \operatorname{tr}(x^2 y^2) \operatorname{tr}(xyxy) + \operatorname{tr}^2(xyxy)) \operatorname{tr}(x^2), \\ v_{15} &= \operatorname{tr}(x^4 \operatorname{tr}(x^2 y) \operatorname{tr}(y^3) + \operatorname{tr}(x^3 y) (-\operatorname{tr}(x^3) \operatorname{tr}(y^3) - 3 \operatorname{tr}(x^2 y) \operatorname{tr}(xy^2)) \\ &\quad + \operatorname{tr}(x^2 y^2) (3 \operatorname{tr}(x^3) \operatorname{tr}(xy^2) + \operatorname{tr}^2(x^2 y)) \\ &\quad + 2 \operatorname{tr}(xyxy) \operatorname{tr}^2(x^2 y) - 4 \operatorname{tr}(xy^3) \operatorname{tr}(x^3) \operatorname{tr}(x^2 y) + \operatorname{tr}(y^4) \operatorname{tr}^2(x^3), \\ v_{16} &= \operatorname{tr}(x^4) \operatorname{tr}^2(xy^2) - 4 \operatorname{tr}(x^3 y) \operatorname{tr}(x^2 y) \operatorname{tr}(xy^2) + \operatorname{tr}(x^2 y^2) (2 \operatorname{tr}(x^3) \operatorname{tr}(xy^2) \\ &\quad + 2 \operatorname{tr}^2(x^2 y)) + 2 \operatorname{tr}(xyxy) \operatorname{tr}^2(x^2 y) - 4 \operatorname{tr}(xy^3) \operatorname{tr}(x^3) \operatorname{tr}(x^2 y) + \operatorname{tr}(y^4) \operatorname{tr}^2(x^3), \\ v_{17} &= \operatorname{tr}(x^2 y^2) (-\operatorname{tr}(x^3) \operatorname{tr}(xy^2) + \operatorname{tr}^2(x^2 y)) + \operatorname{tr}(xyxy) (\operatorname{tr}(x^3) \operatorname{tr}(xy^2) - \operatorname{tr}^2(x^2 y)), \\ v_{18} &= \operatorname{tr}(x^4) (\operatorname{tr}(x^2) \operatorname{tr}(y^2) - \operatorname{tr}^2(xy)) \operatorname{tr}(y^2) \\ &\quad + \operatorname{tr}(x^3 y) (-2 \operatorname{tr}(x^2) \operatorname{tr}(y^2) + 2 \operatorname{tr}^2(xy)) \operatorname{tr}(xy) \\ &\quad + \operatorname{tr}(x^2 y^2) \operatorname{tr}(x^2) (\operatorname{tr}(x^2) \operatorname{tr}(y^2) - \operatorname{tr}^2(xy)), \\ v_{19} &= \operatorname{tr}(x^4) (-\operatorname{tr}(x^2) \operatorname{tr}(y^2) + \operatorname{tr}^2(xy)) \operatorname{tr}(y^2) \\ &\quad + \operatorname{tr}(x^3 y) (2 \operatorname{tr}(x^2) \operatorname{tr}(y^2) - 2 \operatorname{tr}^2(xy)) \operatorname{tr}(xy) \\ &\quad + \operatorname{tr}(xyxy) \operatorname{tr}(x^2) (-\operatorname{tr}(x^2) \operatorname{tr}(y^2) + \operatorname{tr}^2(xy)), \\ v_{20} &= \operatorname{tr}(x^4) \operatorname{tr}(x^2 \operatorname{tr}^2(y^2) - 4 \operatorname{tr}(x^3 y) \operatorname{tr}(x^2) \operatorname{tr}(xy) \operatorname{tr}(y^2) + 4 \operatorname{tr}(x^2 y^2) \operatorname{tr}(x^2) \operatorname{tr}^2(xy) \\ &\quad + 2 \operatorname{tr}(xyxy) \operatorname{tr}^2(x^2) \operatorname{tr}(y^2) - 4 \operatorname{tr}(xy^3) \operatorname{tr}^2(x^2) \operatorname{tr}(xy) + \operatorname{tr}(y^4) \operatorname{tr}^3(x^2), \end{aligned}$$

$$\begin{aligned}
v_{21} &= \text{tr}^2(x^3)\text{tr}^2(y^2) - 4\text{tr}(x^3)\text{tr}(x^2y)\text{tr}(xy)\text{tr}(y^2) \\
&\quad + 2\text{tr}^2(x^2y)\text{tr}(x^2)\text{tr}(y^2) + 4\text{tr}(x^3)\text{tr}(xy^2)\text{tr}^2(xy) + (-2\text{tr}(x^3)\text{tr}(y^3) \\
&\quad - 2\text{tr}(x^2y)\text{tr}(xy^2))\text{tr}(x^2)\text{tr}(xy) + (2\text{tr}(x^2y)\text{tr}(y^3) - \text{tr}^2(xy^2))\text{tr}^2(x^2), \\
v_{22} &= ((\text{tr}(x^3)\text{tr}(xy^2) - \text{tr}^2(x^2y))\text{tr}(y^2) \\
&\quad + (-\text{tr}(x^3)\text{tr}(y^3) + \text{tr}(x^2y)\text{tr}(xy^2))\text{tr}(xy) \\
&\quad + (\text{tr}(x^2y)\text{tr}(y^3) - \text{tr}^2(xy^2))\text{tr}(x^2), \\
v_{23} &= (-\text{tr}(x^3)\text{tr}(xy^2) + \text{tr}^2(x^2y))\text{tr}^2(xy) \\
&\quad + (\text{tr}(x^3)\text{tr}(y^3) - \text{tr}(x^2y)\text{tr}(xy^2))\text{tr}(x^2)\text{tr}(xy) + (-\text{tr}(x^2y)\text{tr}(y^3) + \text{tr}^2(xy^2))\text{tr}^2(x^2), \\
v_{24} &= (\text{tr}(x^2)\text{tr}(y^2) - \text{tr}^2(xy))^2\text{tr}(x^2),
\end{aligned}$$

and

$$\begin{aligned}
\beta_1 &= \frac{1}{4}(\alpha_3 + \alpha_6 - \alpha_7 - \alpha_9), \\
\beta_2 &= \frac{1}{24}(-6\alpha_1 - 3\alpha_2 - 8\alpha_3 - 8\alpha_4 - 17\alpha_5 + 2\alpha_6 - 5\alpha_7 - \alpha_8 - 2\alpha_9 - 6\alpha_{10}), \\
\beta_3 &= \frac{1}{2}(\alpha_3 + \alpha_4 - \alpha_6 + \alpha_7 - 2\alpha_9), \\
\beta_4 &= \frac{1}{12}(-\alpha_2 + \alpha_3 + \alpha_7 - \alpha_9), \\
\beta_5 &= \frac{1}{12}(-12\alpha_1 - 2\alpha_2 + 5\alpha_3 - 3\alpha_4 + 3\alpha_6 + 2\alpha_7 + 4\alpha_9 - 3\alpha_{10}), \\
\beta_6 &= \frac{1}{48}(-3\alpha_2 - 28\alpha_3 - 16\alpha_4 - 37\alpha_5 + 4\alpha_6 - 13\alpha_7 - 5\alpha_8 - 10\alpha_9 - 12\alpha_{10}), \\
\beta_7 &= \frac{1}{48}(-15\alpha_2 - 40\alpha_3 - 16\alpha_4 - 37\alpha_5 + 4\alpha_6 - 13\alpha_7 - 5\alpha_8 - 22\alpha_9 - 24\alpha_{10}), \\
\beta_8 &= \frac{1}{4}(-\alpha_2 + \alpha_3 + 3\alpha_4 + 3\alpha_5 - \alpha_6 - \alpha_7 + \alpha_8 - 3\alpha_{10}), \\
\beta_9 &= \frac{1}{12}(+3\alpha_2 + 5\alpha_3 + 2\alpha_4 + 5\alpha_5 + \alpha_6 + 2\alpha_7 + \alpha_8 - \alpha_9), \\
\beta_{10} &= \frac{1}{12}(-2\alpha_3 + \alpha_4 - 2\alpha_5 + 2\alpha_6 - 2\alpha_7 - 4\alpha_8 + \alpha_9 - 6\alpha_{10}), \\
\beta_{11} &= \frac{1}{8}(+\alpha_2 - \alpha_3 - 2\alpha_4 + \alpha_7 - \alpha_9 + 2\alpha_{10}), \\
\beta_{12} &= \frac{1}{4}(-\alpha_2 + 7\alpha_3 + 2\alpha_4 - 2\alpha_6 + \alpha_7 + 3\alpha_9 + 2\alpha_{10}), \\
\beta_{13} &= \frac{1}{24}(+12\alpha_1 - 6\alpha_2 + 13\alpha_3 + 7\alpha_4 - 14\alpha_5 - 7\alpha_6 - 2\alpha_7 - 10\alpha_8 + 10\alpha_9 + 3\alpha_{10}), \\
\beta_{14} &= \frac{1}{24}(+12\alpha_1 + 6\alpha_2 + 8\alpha_3 + 11\alpha_4 + 17\alpha_5 - 5\alpha_6 + 2\alpha_7 + \alpha_8 + 5\alpha_9 + 9\alpha_{10}), \\
\beta_{15} &= \frac{1}{12}(-\alpha_2 - \alpha_3 - 2\alpha_4 - 8\alpha_5 + 2\alpha_6 - \alpha_7 - 4\alpha_8 + 3\alpha_9), \\
\beta_{16} &= \frac{1}{4}(+\alpha_4 + 3\alpha_5 - \alpha_6 + \alpha_8 - \alpha_9 - \alpha_{10}), \\
\beta_{17} &= \frac{1}{12}(+\alpha_2 - 2\alpha_3 - \alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7 - 2\alpha_8 - 3\alpha_9 - 3\alpha_{10}), \\
\beta_{18} &= \frac{1}{48}(-15\alpha_2 - 64\alpha_3 - 22\alpha_4 - 43\alpha_5 + 10\alpha_6 - 19\alpha_7 - 11\alpha_8 - 28\alpha_9 - 36\alpha_{10}), \\
\beta_{19} &= \frac{1}{48}(-9\alpha_2 - 46\alpha_3 - 22\alpha_4 - 43\alpha_5 + 10\alpha_6 - 13\alpha_7 - 11\alpha_8 - 22\alpha_9 - 24\alpha_{10}),
\end{aligned}$$

$$\begin{aligned}
\beta_{20} &= \frac{1}{96}(+9\alpha_2 + 4\alpha_3 + 10\alpha_4 + 25\alpha_5 + 2\alpha_6 + \alpha_7 + 5\alpha_8 + 4\alpha_9), \\
\beta_{21} &= \frac{1}{48}(+3\alpha_2 + 9\alpha_3 - \alpha_4 + 3\alpha_5 - \alpha_6 + 5\alpha_7 + 5\alpha_8 + 11\alpha_{10}), \\
\beta_{22} &= \frac{1}{48}(-3\alpha_2 + 18\alpha_3 + 4\alpha_4 + 15\alpha_5 - 14\alpha_6 + 7\alpha_7 + 7\alpha_8 + 10\alpha_{10}), \\
\beta_{23} &= \frac{1}{4}(+2\alpha_3 + \alpha_5 - \alpha_6 + \alpha_7 + \alpha_8 + 2\alpha_{10}), \\
\beta_{24} &= \frac{1}{16}(+\alpha_2 + 2\alpha_3 + \alpha_5 + \alpha_7 + \alpha_8 + 2\alpha_{10}).
\end{aligned}$$

Lemma 3.9. For $\lambda = (5, 5)$ in C_0 :

$$\begin{aligned}
w_1 - 5v_2 &= 0, \\
w_2 - w_3 + v_1 - 4v_2 - v_3 &= 0, \\
w_3 + w_4 - \frac{5}{2}v_1 - v_2 + 2v_3 &= 0,
\end{aligned}$$

where

$$\begin{aligned}
v_1 &= \text{tr}([x, y]^3 x^2) \text{tr}(y^2) - (\text{tr}([x, y]^3 yx) + \text{tr}([x, y]^3 xy) \text{tr}(xy) + \text{tr}([x, y]^3 y^2) \text{tr}(x^2)), \\
v_2 &= \frac{1}{2}(\text{tr}(x^2 y^2 xy) - \text{tr}(y^2 x^2 yx)) \text{tr}([x, y]^2), \\
v_3 &= (\text{tr}(x^2 y^2 xy) - \text{tr}(y^2 x^2 yx))(\text{tr}(x^2) \text{tr}(y^2) - \text{tr}^2(xy)), \\
v_4 &= (\text{tr}(x^3) \text{tr}(y^2) - 2\text{tr}(x^2 y) \text{tr}(xy) + \text{tr}(xy^2) \text{tr}(x^2)) \text{tr}([x, y]^2 y) \\
&\quad - (\text{tr}(y^3) \text{tr}(x^2) - 2\text{tr}(xy^2) \text{tr}(xy) + \text{tr}(x^2 y) \text{tr}(y^2)) \text{tr}([x, y]^2 x).
\end{aligned}$$

The following theorem is the main result of the paper.

Theorem 3.10. The minimal generating GL_2 -module G of the pure trace algebra C_{42} of two generic 4×4 matrices decomposes as

$$\begin{aligned}
G &= W(1, 0) \oplus W(2, 0) \oplus W(3, 0) \oplus W(4, 0) \oplus W(2, 2) \oplus W(3, 2) \\
&\oplus W(4, 2) \oplus W(3, 3) \oplus W(4, 3) \oplus W(5, 3) \oplus W(4, 4) \oplus W(6, 3) \oplus W(5, 5).
\end{aligned}$$

For all $\lambda = (\lambda_1, \lambda_2) \neq (5, 5)$, the module $W(\lambda_1, \lambda_2)$ is generated by the element

$$w_\lambda(X, Y) = \text{tr}((XY - YX)^{\lambda_2} X^{\lambda_1 - \lambda_2}).$$

For $\lambda = (5, 5)$ a generator of $W(5, 5)$ is

$$w_{(5,5)}(X, Y) = \text{tr}((XY - YX)^3 (X^2 Y^2 - X Y Y X - Y X X Y + Y^2 X^2)).$$

Proof. Using (5), we may replace the elements $\text{tr}(Z_1 \cdots Z_k)$, $Z_i = X, Y$, $k \geq 2$, modulo the subalgebra generated by products of traces of degree $< k$, with $\text{tr}(z_1 \cdots z_k)$, where $z_i = x, y$ are generic traceless matrices. Hence, it is sufficient to show that the algebra C_0 has a minimal GL_2 -module of generators

$$\begin{aligned}
(9) \quad M &= W(2, 0) \oplus W(3, 0) \oplus W(4, 0) \oplus W(2, 2) \oplus W(3, 2) \\
&\oplus W(4, 2) \oplus W(3, 3) \oplus W(4, 3) \oplus W(5, 3) \oplus W(4, 4) \oplus W(6, 3) \oplus W(5, 5),
\end{aligned}$$

each $W(\lambda_1, \lambda_2)$, $(\lambda_1, \lambda_2) \neq (5, 5)$, is generated by

$$w_\lambda(x, y) = \text{tr}((xy - yx)^{\lambda_2} x^{\lambda_1 - \lambda_2}),$$

and for $\lambda = (5, 5)$ a generator of $W(5, 5)$ is

$$(10) \quad w_{(5,5)}(x, y) = \text{tr}((xy - yx)^3 (x^2 y^2 - xy yx - yx xy + y^2 x^2)).$$

There are three typical cases. We consider the image of the GL_2 -modules U_n in C_0 under the natural homomorphism $\pi : P(x, y) \rightarrow C_0$. We may assume that the generators of C_0 are in $\pi(U_2 \oplus \cdots \oplus U_{10})$. Lemma 3.1 allows to consider the cases $n \geq 6$ only.

Let $\lambda = (n)$, $n = 6, \dots, 10$. This case was handled in Lemma 3.1. The only GL_2 -module $W(n)$ in $\pi(U_n)$ is generated by $\text{tr}(x^n)$. By (7), $\text{tr}(x^5)$ belongs to the subalgebra of C_0 generated by traces of lower degree. Similar equations hold for the other $n \geq 6$. Hence, $W(n)$ does not participate in any minimal GL_2 -module of generators of C_0 for $n = 6, \dots, 10$.

Let $\lambda = (6, 2)$. By Proposition 2.1, $W(6, 2)$ participates with multiplicity 3 in U_8 . The highest weight vectors of the three copies of $W(6, 2)$ are w_1, w_2, w_3 from Lemma 2.2. By the relations in Lemma 3.4, $\pi(w_1), \pi(w_2), \pi(w_3)$ are linear combinations of $v_1, \dots, v_{10}$, and hence belong to the subalgebra of C_0 generated by the elements of degree ≤ 7 (and even of degree ≤ 6). Therefore, as in the case $\lambda = (n)$, the minimal GL_2 -modules of generators of C_0 do not contain $W(6, 2)$. The cases $\lambda = (5, 2), (7, 2), (5, 4), (8, 2), (7, 3), (6, 4)$ are similar.

Let $\lambda = (5, 3)$. By Proposition 2.1 again, $W(5, 3)$ participates with multiplicity 3 in U_8 . The corresponding highest weight vectors are w_1, w_2, w_3 from Lemma 2.2, and the element w_1 is equal to $\text{tr}((xy - yx)^3 x^2)$. By the relations in Lemma 3.4, $\pi(w_2), \pi(w_3)$ are linear combinations of $\pi(w_1)$ and $v_1, \dots, v_5$. Hence they belong to the subalgebra of C_0 generated by the elements of degree ≤ 7 and by the GL_2 -module generated by w_1 . In this way, the multiplicity of $W(5, 3)$ in the minimal GL_2 -module of generators is bounded from above by 1. The cases $\lambda = (4, 2), (4, 3), (4, 4), (6, 3)$ are similar. The case $\lambda = (5, 5)$ is analogous but one has to express the elements w_1, w_3, w_4 in terms of w_2 and v_1, v_2, v_3, v_4 . It is easy to see that w_2 is equal to (10). Finally, the multiplicity of $W(3, 3)$ in U_6 is already 1. In this way we obtain that the minimal GL_2 -module of generators of C_0 is a homomorphism image of the module M from (9), and the algebra C_0 is a homomorphic image of the symmetric algebra $K[M]$ of M .

Clearly, the coefficients of the Hilbert series of $K[M]$ are bigger or equal to the coefficients of the Hilbert series of C_0 . The Hilbert series of C_{42} is given in (1). The factor $(1 - t)(1 - u)$ in its denominator corresponds to the polynomial algebra $K[\text{tr}(X), \text{tr}(Y)]$ in (6). Hence, the Hilbert series of C_0 is equal to $h(t, u) = P_C(t, u)/Q_C(t, u)$ from (2). It follows from (9) that the Hilbert series of $K[M]$ is

$$H(K[M], t, u) = \frac{1}{q_2(t, u) \cdots q_{10}(t, u)},$$

where $q_2(t, u), \dots, q_5(t, u)$ are given in (8) and

$$q_6(t, u) = (1 - t^4 u^2)(1 - t^3 u^3)^2(1 - t^2 u^4),$$

$$q_7(t, u) = (1 - t^4 u^3)(1 - t^3 u^4),$$

$$q_8(t, u) = (1 - t^5 u^3)(1 - t^4 u^4)^2(1 - t^3 u^5),$$

$$q_9(t, u) = (1 - t^6 u^3)(1 - t^5 u^4)(1 - t^4 u^5)(1 - t^3 u^6), \quad q_{10}(t, u) = (1 - t^5 u^5).$$

By direct calculations we see that the coefficients of the homogeneous components of degree ≤ 10 of $H(K[M], t, u)$ are the same as the coefficients $h_0, h_1, \dots, h_{10}$ from (4). This implies that the generators (9) satisfy no relations of degree ≤ 10 and completes the proof of the theorem. $\square$

Remark 3.11. Since $\text{tr}([X, Y]) = 0$, the equation (7) gives that the element $\text{tr}([X, Y]^5)$ of C_{42} belongs to the subalgebra generated by traces of lower degree.

Remark 3.12. Calculating the first homogeneous components of the difference $H(C_0, t, u) - H(K[M], t, u)$, we see that

$$\begin{aligned} H(C_0, t, u) - H(K[M], t, u) = & (S_{(7,5)}(t, u) + 2S_{(6,6)}(t, u)) + (S_{(8,5)}(t, u) + 2S_{(7,6)}(t, u)) \\ & + (2S_{(9,5)}(t, u) + 6S_{(8,6)}(t, u) + 2S_{(7,7)}(t, u)) \\ & + (2S_{(10,5)}(t, u) + 9S_{(9,6)}(t, u) + 7S_{(8,7)}(t, u)). \end{aligned}$$

Hence, the GL_2 -modules of the defining relations of degree 12 and 13 are, respectively, $W(7, 5) \oplus 2W(6, 6)$ and $W(8, 5) \oplus 2W(7, 6)$. In principle, the defining relations can be found as those in Lemmas 3.2 – 3.9, but the calculations should be more complicated.

Remark 3.13. The homogeneous system of parameters of C_{42} found by Teranishi [T1, T2] contains all traces of degree ≤ 4 and two elements of degree $(4, 2)$ and $(2, 4)$, respectively. Translated in our language, we can choose for a homogeneous system of parameters of C_{42} the elements

$$\begin{aligned} & \text{tr}(X), \text{tr}(Y), \text{tr}(X^2), \text{tr}(XY), \text{tr}(Y^2), \\ & \text{tr}(X^3), \text{tr}(X^2Y), \text{tr}(XY^2), \text{tr}(Y^3), \\ & \text{tr}(X^4), \text{tr}(X^3Y), \text{tr}(X^2Y^2), \text{tr}(XYXY), \text{tr}(XY^3), \text{tr}(Y^4), \end{aligned}$$

and the polynomials

$$w_{(4,2)}(X, Y) = \text{tr}((XY - YX)^2 X^2), \quad w_{(2,4)}(Y, X) = \text{tr}((XY - YX)^2 Y^2)$$

from Theorem 3.10.

REFERENCES

- [AP] S. Abeasis, M. Pittaluga, *On a minimal set of generators for the invariants of 3×3 matrices*, Commun. Algebra **17** (1989), 487-499.
- [ADF] G. Almkvist, W. Dicks, E. Formanek, *Hilbert series of fixed free algebras and noncommutative classical invariant theory*, J. Algebra **93** (1985), 189-214.
- [ADS] H. Aslaksen, V. Drensky, L. Sadikova, *Defining relations of invariants of two 3×3 matrices*, J. Algebra **298** (2006), 41-57.
- [BS] A. Berele, J.R. Stembridge, *Denominators for the Poincaré series of invariants of small matrices*, Israel J. Math. **114** (1999), 157-175.
- [DEP] C. De Concini, D. Eisenbud, C. Procesi, *Young diagrams and determinantal varieties*, Invent. Math. **56** (1980), 129-165.
- [D1] V. Drensky, *Free Algebras and PI-Algebras*, Springer-Verlag, Singapore, 1999.
- [D2] V. Drensky, *Defining relations for the algebra of invariants of 2×2 matrices*, Algebras and Representation Theory **6** (2003), 193-214.
- [DF] V. Drensky, E. Formanek, *Polynomial Identity Rings*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser Verlag, Basel, 2004.
- [Du] J. Dubnov, *Sur une généralisation de l'équation de Hamilton-Cayley et sur les invariants simultanés de plusieurs affineurs*, Proc. Seminar on Vector and Tensor Analysis, Mechanics Research Inst., Moscow State Univ. **2/3** (1935), 351-367 (see also Zbl. für Math. **12** (1935), p. 176).
- [F] E. Formanek, *The Polynomial Identities and Invariants of $n \times n$ Matrices*, CBMS Regional Conf. Series in Math. **78**, Published for the Confer. Board of the Math. Sci. Washington DC, AMS, Providence RI, 1991.
- [K] P.E. Koshlukov, *Polynomial identities for a family of simple Jordan algebras*, Commun. Algebra **16** (1988), 1325-1371.

- [Ku] E.N. Kuzmin, *On the Nagata-Higman theorem* (Russian), in “Mathematical Structures, Computational Mathematics, Mathematical Modelling. Proc. Dedicated to the 60th Birthday of Acad. L. Iliev”, Sofia, 1975, 101-107.
- [N] K. Nakamoto, *The structure of the invariant ring of two matrices of degree 3*, J. Pure Appl. Algebra **166** (2002), No. 1-2, 125-148.
- [R] Yu.P. Razmyslov, *Trace identities of full matrix algebras over a field of characteristic zero* (Russian), Izv. Akad. Nauk SSSR, Ser. Mat. **38** (1974), 723-756. Translation: Math. USSR, Izv. **8** (1974), 727-760.
- [SZ] I. Shestakov, N. Zhukavets, On associative algebras satisfying the identity $x^5 = 0$, Algebra Discrete Math. (Lugansk) (2004), No. 1, 112-120.
- [S] K.S. Sibirskii, *Algebraic invariants for a set of matrices* (Russian), Sib. Mat. Zh. **9** (1968), No. 1, 152-164. Translation: Siber. Math. J. **9** (1968), 115-124.
- [T1] Y. Teranishi, *The ring of invariants of matrices*, Nagoya Math. J. **104** (1986), 149-161.
- [T2] Y. Teranishi, *Linear Diophantine equations and invariant theory of matrices*, Commutative algebra and combinatorics (Kyoto, 1985), 259-275, Adv. Stud. Pure Math., **11**, North-Holland, Amsterdam, 1987.
- [V] M. Van den Bergh, *Explicit rational forms for the Poincaré series of the trace rings of generic matrices*, Isr. J. Math. **73** (1991), 17-31.
- [VL] M. R. Vaughan-Lee, *An algorithm for computing graded algebras*, J. Symbolic Comput. **16** (1993), 345-354.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, 1113
SOFIA, BULGARIA

E-mail address: `drensky@math.bas.bg`

FACHGRUPPE INFORMATIK, RWTH AACHEN, 52056 AACHEN, GERMANY

E-mail address: `sadikova@stce.rwth-aachen.de`